

Effect of Diabetes Self Management Education on Blood Sugar Level Control for Diabetics in Padangsidempuan City

Adi Antoni¹, Eva Decroli², Afriwardi³, Irwan Prayitno⁴, Nur Indrawati Lipoeto⁵, Nursyirwan Efendi⁶, Mudjiran⁷, Hardisman⁸

Abstract

Diabetes mellitus (DM) is a chronic disease whose number is always increasing every year in the world. DM requires serious treatment in order to reduce the complications of this disease. The purpose of this study was to determine the effect of diabetes self-management education on controlling blood sugar levels in diabetics in the city of Padangsidempuan. This study used a one group pretest-posttest only design involving 40 respondents based on sample size calculation using sample size determination in health studies. The sample technique uses proportional random sampling. The sample criteria are type 2 diabetes mellitus, diabetic patients who do not experience cognitive impairment, can communicate verbally and nonverbally. Measuring blood sugar levels using HbA1c. The results of this study indicate that the average before the DSME intervention obtained an HbA1c value of 8.828% (SD=.17725) and the average after the DSME intervention obtained an HbA1c value of 7.940% (1.1502) (p value = <0.001). DSME is effective in reducing HbA1c values in people with diabetes mellitus. So it is hoped that DM sufferers can take advantage of DSME as an independent intervention that can be done at home to prevent various problems from DM.

Keywords: diabetes self management education, diabetes mellitus, blood sugar levels, HbA1c.

Introduction

Diabetes mellitus is an epidemic disease throughout the world with an ever-increasing prevalence and is a global health burden. It is estimated that 537 million people have diabetes at the age of 29-79 years and it is predicted to increase to 783 million people in 2045(1). Approximately 422 million people worldwide have diabetes, the majority of whom live in low- and middle-income countries, and 1.6 million deaths are directly attributed to diabetes each year (2). There are 536.6 million people suffering from diabetes in the world in 2021 and 205.6 million people in the WP (Western Pacific) Region and will increase in 2045 to 260.2 million (3).

¹ Student of Doctoral Program of Public Health Sciences, Faculty of Medicine, University of Andalas, Padang, Indonesia, e-mail: adiantoni100@gmail.com

² Faculty of Medicine, Andalas University, Padang, West Sumatra, Indonesia

³ Faculty of Medicine, Andalas University, Padang, West Sumatra, Indonesia

⁴ Universitas Muhammadiyah Jakarta, Jakarta, Indonesia

⁵ Faculty of Medicine, Andalas University, Padang, West Sumatra, Indonesia

⁶ Department of Anthropology, Andalas University, Padang, West Sumatra, Indonesia

⁷ Guidance and Counselling Department, Universitas Negeri Padang, West Sumatera, Indonesia

⁸ Faculty of Medicine, Andalas University, Padang, West Sumatra, Indonesia

The total cases of diabetes in Indonesia are 19.5 million people in 2021 and will increase to 28.6 million people in 2045 while Indonesia is ranked 5th in the world with the most diabetes sufferers, namely 10.2 million people and in 2021(4). The 2018 Riskesdas results found an increase in the prevalence of people with diabetes mellitus (age > 15 years) from 6.9% in 2013 to 10.9% in 2018 (5). North Sumatra is one of the provinces with the fourth most diabetes sufferers in Indonesia. Types of diabetes mellitus control in North Sumatra based on eating arrangements were obtained at 78.3%, exercise at 46.1%, and herbal alternatives at 37.1%(6).

Controlled blood sugar levels are one measure of the success of self-management of people with diabetes mellitus, however, the target for achieving glycemetic control is still not satisfactory and most of them are still above the desired target of 7% (7). One of the causes of low glycemetic control is low adherence to taking medication and long duration of illness affects the low glycemetic control of DM patients (8)

Higher diabetes knowledge and higher medication adherence were significant predictors of good glycemetic control (9). There is a negative correlation between knowledge about diabetes and HbA1c values. This means that the higher the knowledge about diabetes, the lower the HbA1c value. In addition, there is a positive correlation between knowledge about diabetes and self-care management (10). In addition, DSME can increase the chances of glycemetic control with an estimated value before intervention of 0.53 to 0.82 (14).

Based on the description above, researchers are interested in researching the effect of diabetes self-management education on controlling blood sugar levels of diabetics in Padangsidempuan City.

Method

This study used a Cluster Randomized Controlled Trial (CRCT) based on the CONSORT (Consolidated Standards of Reporting Trials) guidelines. The design of this study was a quasy experiment with a one group pre test post test only design which was carried out in the city of Padangsidempuan. The number of samples in this study were 40 respondents using the Sample Size Determination in Health Studies formula. The sample technique uses proportional random sampling. The sample criteria in this study were type 2 diabetes mellitus, diabetic patients who did not experience cognitive impairment, were able to communicate verbally and nonverbally.

Measurement of blood sugar levels using Glycated hemoglobin (HbA1c). Measured using HPLC (High Performance Liquid Chromatography). This research was conducted for 3 months. Data analysis using the Paired t Test

Results

Table 1. Average HBA1c values before and after the DSME educational intervention in diabetes mellitus patients (n = 40)

Variable	Before intervention (n=40)			After Intervention (n=40)			p value
	Mean	SD	95%CI	Mean	SD	95%CI	
HBA1c	8.828	1.17725	8.453 - 9.202	7.940	1.1502	7.572 - 8.308	<0.001

The table above shows the results of the paired t test on HBA1c in the first measurement (pretest) obtained an average HBA1c value of 8.828% while the average HBA1c after the intervention was 7.940%. the difference in HBA1c before and after the intervention was 0.888%. Based on the results of the statistical test, it can be concluded that there is an effect of diabetes self-management on blood sugar levels in people with diabetes mellitus.

Discussion

The results of this study indicate that DSME is able to control blood sugar levels. This is in accordance with previous studies, namely there is a decrease in HbA1c values (15). Chai stated that out of 118 respondents (63 intervention groups, 55 control groups) DSME could significantly reduce anxiety, depression, and HbA1c scores (before 7.2%, after 6.2%), blood sugar levels 2 hours after the tolerance test oral glucose or OGTT and fasting blood sugar (16).

DSME can reduce HbA1c values (before 10.51% and after 9.89%) (14). In addition, DSME can lower HbA1c values in the absence of hypoglycemia (17). DSME also affects physical activity and dietary habits of people with diabetes mellitus (18).

Adequate glycemic control or HbA1c is very difficult for diabetics (19). Diabetes self-management is influenced by four factors, namely self-efficacy, level of knowledge, spiritual level and social support (20). Several studies have shown that education in diabetes patients that refers to self-efficacy is very effective in improving self-care (21)(22)(23).

Diabetics who did not receive education about diabetes had a knowledge score of 15.3% lower than those who received education and had an average HbA1c value of 0.89% higher than those who received education (24). In addition, poor disease knowledge and non-adherence to treatment among DM patients has been an ongoing problem for both patients and healthcare providers (25).

Educational programs that emphasize lifestyle modification with the importance of adherence to the overall medication regimen, especially diet, exercise, and regular follow-up will be more beneficial in glycemic control than adherence to medication alone (26) (27).

Management of diabetes mellitus is very important in preventing the dangers of complications as well as reducing the number of people with diabetes and reducing the cost of expensive treatment (28). In general, the management of diabetes mellitus aims to improve the quality of life. Management of diabetes mellitus specifically in the form of education, medical nutritional therapy, physical exercise, and pharmacological therapy (7).

Diabetes self-management is at the heart of diabetes management (29). Diabetics need about 99% self-care independently in managing their disease (30). Things that are needed in the self-management of diabetes patients are proper education, lifestyle modification, medication and blood glucose control. Education or education is the foundation in achieving the success of diabetes management (31).

In order to improve the health status of the people in Indonesia and the success of the social security program in the health sector, in accordance with Presidential Regulation Number 12 of 2013 concerning Health Insurance Article 21 Paragraph 1, one of Prolanis' activities through group education is a form of promotive and preventive efforts related to chronic diseases experienced by people, especially diabetes. An indicator of the success of implementing Prolanis in a health center is that 75% of participants have "good" results on specific examinations for diabetes mellitus according to clinical guidelines (32).

Even though the implementation of the educational program has been implemented for a long time, there are still deficiencies in its application. Based on the research by Raraswati, Keryawan and Soetedjo in 2018, it was found that the implementation of education in the prolanis program at the Jatinangor District Health Center had been carried out but was not optimal. This is caused by several factors, among others. The doctor's factor that might occur is the lack of education of the doctor to the patient about the disease he is suffering from and it is important to carry out monitoring so that the

patient's awareness to check fasting blood sugar every month is low. The patient factor, that most patients only come to the health center when there are complaints or just to take medicine do not do routine fasting blood sugar checks (33).

Conclusions and recommendations

Based on this study it can be concluded that diabetes education through DMSE can be a way to control blood sugar levels in patients with diabetes mellitus. So that the blood sugar levels of diabetics remain stable. Because diabetes is a chronic disease that requires lifelong intervention.

The results of this study are expected to be one of the independent interventions for people with diabetes mellitus at home.

References

1. International Diabetes Federation. International Diabetes Federation Atlas 10th Edition. Vol. 102, Diabetes Research and Clinical Practice. 2021. 147–148 p.
2. National Diabetes Statistics. National Diabetes Statistics Report, 2020. Natl Diabetes Stat Rep. 2020;2.
3. IDF. International Diabetes Federation: Diabetes Atlas. 2020.
4. Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract* [Internet]. 2018;138:271–81. Available from: <https://doi.org/10.1016/j.diabres.2018.02.023>
5. KEMENKES. Hasil Utama Riskesdas Tentang Prevalensi Diabetes Mellitus di Indonesia 2018. Hasil Utama Riskesdas Tentang Prevalensi Diabetes Mellitus di Indonesia 2018. 2018;8.
6. Riskesdas. Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI. *J Chem Inf Model* [Internet]. 2018;53(9):1689–99. Available from: http://www.yankes.kemkes.go.id/assets/downloads/PMK_No_57_Tahun_2013_tentang_PTRM.pdf
7. PERKENI. Pedoman Pengelolaan dan Pencegahan Diabetes Mellitus Tipe 2 Dewasa di Indonesia 2021. *Glob Iniat Asthma* [Internet]. 2021;46. Available from: www.ginasthma.org.
8. McElfish PA, Riklon S, Purvis RS, Long CR, Felix HC, Hudson JS, et al. Study protocol for family model diabetes self-management education with Marshallese participants in faith-based organizations. *Contemp Clin Trials Commun* [Internet]. 2022;30(September):101007. Available from: <https://doi.org/10.1016/j.conctc.2022.101007>
9. Al-Qazaz HK, Sulaiman SA, Hassali MA, Shafie AA, Sundram S, Al-Nuri R, et al. Diabetes knowledge, medication adherence and glycemic control among patients with type 2 diabetes. *Int J Clin Pharm*. 2011;33(6):1028–35.
10. Bukhsh A, Khan TM, Nawaz MS, Ahmed HS, Chan KG, Goh BH. Association of diabetes knowledge with glycemic control and self-care practices among Pakistani people with type 2 diabetes mellitus. *Diabetes, Metab Syndr Obes Targets Ther*. 2019;12:1409–17.
11. Haas L, Maryniuk M, Beck J, Cox CE, Duker P, Edwards L, et al. National Standards for Diabetes Self-Management Education and Support. *Diabetes Educ*. 2012;38(5):619–29.
12. ADA. Diagnosis and Classification of Diabetes Mellitus DEFINITION AND DESCRIPTION OF DIABETES MELLITUS. *Diabetes Care*. 2014;37(January):81–90.
13. Wichit N, Mnatzaganian G, Courtney M, Schulz P, Johnson M. Randomized controlled trial of a family-oriented self-management program to improve self-efficacy, glycemic control and quality of life among Thai individuals with Type 2 diabetes. *Diabetes Res Clin Pract* [Internet]. 2017;123:37–48. Available from: <http://dx.doi.org/10.1016/j.diabres.2016.11.013>

14. McElfish PA, Long CR, Kohler PO, Yeary KHK, Bursac Z, Narcisse MR, et al. Comparative effectiveness and maintenance of diabetes self-management education interventions for Marshallese patients with type 2 diabetes: A randomized controlled trial. *Diabetes Care*. 2019;42(5):849–58.
15. Adam L, O'Connor C, Garcia AC. Evaluating the Impact of Diabetes Self-Management Education Methods on Knowledge, Attitudes and Behaviours of Adult Patients With Type 2 Diabetes Mellitus. *Can J Diabetes* [Internet]. 2018;42(5):470-477.e2. Available from: <https://doi.org/10.1016/j.jcjd.2017.11.003>
16. Chai S, Yao B, Xu L, Wang D, Sun J, Yuan N, et al. The effect of diabetes self-management education on psychological status and blood glucose in newly diagnosed patients with diabetes type 2. *Patient Educ Couns* [Internet]. 2018;101(8):1427–32. Available from: <https://doi.org/10.1016/j.pec.2018.03.020>
17. Boels AM, Vos RC, Dijkhorst-Oei LT, Rutten GEHM. Effectiveness of diabetes self-management education and support via a smartphone application in insulin-treated patients with type 2 diabetes: Results of a randomized controlled trial (TRIGGER study). *BMJ Open Diabetes Res Care*. 2019;7(1):1–10.
18. Abdulah DM, Hassan AB, Saadi FS, Mohammed AH. Impacts of self-management education on glycaemic control in patients with type 2 diabetes mellitus. *Diabetes Metab Syndr Clin Res Rev* [Internet]. 2018;12(6):969–75. Available from: <https://doi.org/10.1016/j.dsx.2018.06.007>
19. Gamboa Moreno E, Mateo-Abad M, Ochoa de Retana García L, Vrotsou K, del Campo Pena E, Sánchez Perez Á, et al. Efficacy of a self-management education programme on patients with type 2 diabetes in primary care: A randomised controlled trial. *Prim Care Diabetes*. 2019;13(2):122–33.
20. Sonsona JB. Factors Influencing Diabetes Self-Management of Filipino Americans with Type 2 Diabetes Mellitus: A Holistic Approach. *ProQuest Diss Theses* [Internet]. 2014;178. Available from: https://search.proquest.com/docview/1513243781?accountid=28931%0Ahttp://metaiskalnik.i zum.si:80/sfxlcl41?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dissertation&genre=dissertations+%26+theses&sid=ProQ:ProQuest+Dissertations+%26+Theses+Global&ati
21. Lorig K, Ritter PL, Villa FJ, Armas J. Community-based peer-led diabetes self-management: A randomized trial. *Diabetes Educ*. 2009;35(4):641–51.
22. King DK, Glasgow RE, Toobert DJ, Strycker LA, Estabrooks PA, Osuna D, et al. Self-efficacy, problem solving, and social-environmental support are associated with diabetes self-management behaviors. *Diabetes Care*. 2010;33(4):751–3.
23. Yoo H, Kim CJ, Jang Y, You MA. Self-efficacy associated with self-management behaviours and health status of South Koreans with chronic diseases. *Int J Nurs Pract*. 2011;17(6):599–606.
24. Phillips E, Rahman R, Mattfeldt-Beman M. Relationship between diabetes knowledge, glycemic control, and associated health conditions. *Diabetes Spectr*. 2018;31(2):196–9.
25. Morgan C, Currie C, Peters J. Relationship Between Diabetes and Mortality. *Diabetes Care*. 2000;23(8):1103–7.
26. Khattab M, Khader YS, Al-Khawaldeh A, Ajlouni K. Factors associated with poor glycemic control among patients with Type 2 diabetes. *J Diabetes Complications* [Internet]. 2010;24(2):84–9. Available from: <http://dx.doi.org/10.1016/j.jdiacomp.2008.12.008>
27. Al-Rasheedi AAS. The Role of Educational Level in Glycemic Control among Patients with Type II Diabetes Mellitus. *Int J Health Sci (Qassim)*. 2014;8(2):177–87.
28. Schinckus L, Van den Broucke S, Housiaux M. Assessment of implementation fidelity in diabetes self-management education programs: A systematic review. *Patient Educ Couns*. 2014;96(1):13–21.

29. Ausili D, Barbaranelli C, Rossi E, Rebora P, Fabrizi D, Coghi C, et al. Development and psychometric testing of a theory-based tool to measure self-care in diabetes patients: The Self-Care of Diabetes Inventory. *BMC Endocr Disord.* 2017;17(1):1–12.
30. Funnell MM, Anderson RM. Empowerment and Self-Management of Diabetes. *Clin Diabetes.* 2004;22(3):123–7.
31. Haas L, Maryniuk M, Beck J, Cox CE, Duker P, Edwards L, et al. Professional practice committee for the 2014 clinical practice recommendations. *Diabetes Care.* 2014;37(SUPPL.1):1630–7.
32. BPJS Kesehatan. Panduan praktis Prolanis (Program pengelolaan penyakit kronis). BPJS Kesehatan. 2014;
33. Raraswati A, Heryaman H, Soetedjo NNM. Peran Program Prolanis dalam Penurunan Kadar Gula Darah Puasa pada Pasien Diabetes Melitus Tipe 2 di Puskesmas Kecamatan Jatinangor. *J Sist Kesehat [Internet].* 2018;4(2):65–70. Available from: http://jurnal.unpad.ac.id/jsk_ikm/article/view/20687