

Detection Of Pathogens In Human Consumption Water Through Pcr Technique

Darwin Núñez^{1,2*}, Maria Monserrath Morales², Diego Moposita³, Jose Luis Altuna⁴

ABSTRACT

In this study, the presence of pathogenic microorganisms such as Listeria Monocytogenes, Escherichia coli with its O157:H7 serotype, and Salmonella spp. was determined in drinking water in the Chambo canton. Selective culture media techniques and conventional Polymerase Chain Reaction (PCR) were employed for this purpose. Specific primers for each pathogen were used for thermocycling, along with positive controls for contrast. A total of 50 samples from different strategic points in the area were analyzed using these methodologies. Listeria monocytogenes was detected in 40% (20/50) of the samples, Escherichia coli with its O157:H7 serotype in 80% (40/50), and Salmonella spp. in 20% (10/50). Thus, the quality of the drinking water in the canton was determined based on the number of points contaminated with the studied pathogens. The objective of this research is to understand the degree of contamination in drinking water and to consider possible alternatives in the sector to reduce the microbial load in the study matrix.

KEYWORD: PCR, Water Quality, Pathogens.

OBJECTIVE

The objective of this research is to determine the quality of water for human consumption in the Chambo canton, using molecular techniques that provide a reliable result and future decision-making in the face of possible threats.

INTRODUCCION

Water is the fundamental element for the development of vital activities on planet Earth. This life-giving liquid is present in an approximate quantity of 1.4 trillion cubic kilometers. It's important to note that the majority of this water is saline and is primarily found in the oceans and seas, accounting for 97.5% of the total ⁽¹⁾.

Freshwater, which is essential for human consumption and terrestrial activities, makes up approximately 2.5% of the total water on Earth ⁽²⁾. Of this freshwater, the majority exists in the form of ice in polar ice caps, glaciers, and ice sheets, accounting for approximately 68.7%. Groundwater, found in aquifers and underground reservoirs, makes up about 30.1%. The remaining freshwater can be found in lakes, rivers, soil moisture, and in the atmosphere ⁽³⁾.

In the Chambo canton of the Chimborazo province, water is used for both human consumption and irrigation. However, the conditions from the water sources to reservoirs and distribution are not adequate. This is because as the water flows through channels, streams, and springs, it is prone to contamination by animals such as cows, horses, and animals in a state of decomposition. These animals approach the channels to consume and defecate, leading¹ to contamination and the proliferation of pathogenic microorganisms.

¹ Universidad Estatal de Bolívar, Facultad Ciencias Agropecuarias, Carrera Agroindustrias
<https://orcid.org/0000-0003-0338-3759>

² Universidad Técnica de Cotopaxi, Unidad Académica de Posgrado, Ecuador

This contamination can have a severe impact on the health of those who consume this water, causing serious injuries or even death.

Detecting pathogens in drinking water is of great importance to ensure safety and public health ⁽⁴⁾. The presence of pathogenic microorganisms in water intended for human consumption can lead to waterborne disease outbreaks and pose a significant risk to the population ⁽⁵⁾.

The molecular biology technique known as Polymerase Chain Reaction (PCR) is a widely used and effective tool for detecting pathogens in samples of water for human consumption. PCR allows for the selective amplification and detection of specific DNA fragments from pathogens, even at low concentrations ⁽⁶⁾.

Hence, the general objective of the current research was to detect pathogens in water for human consumption using the PCR technique. By conducting analyses through PCR, reliable data can be obtained to make accurate assessments of the degree of pathogenic contamination that may be present in drinking water. This information can, in turn, guide appropriate corrective actions in the event of biological contaminants being detected.

MATERIALS AND METHODS

Study Area: The present study is descriptive, experimental, and cross-sectional in nature. It was conducted from January to September 2023, focusing on the probability of the presence of pathogens such as *Salmonella* spp., *Listeria* spp., *E. coli*, and *Listeria* in 50 samples of drinking water. These samples were collected from 50 strategic points, with 10 samples from each of the North, South, East, West, and Central areas of the Chambo canton, ranging from the sources of human drinking water to households ⁽⁷⁾.

Georeferencing: This method allowed for the precise or relative location of different sampling points. Coordinates were determined using the GPSMAP 64sx, which provided the necessary data to be processed in the ARCGIS program, resulting in a digital map ⁽⁸⁾.

Sampling Technique: The collection of samples of drinking water was performed using sterile polyethylene food-grade materials, properly labeled. Strict cold chain procedures were maintained during transportation, and samples were isolated from natural elements ⁽⁹⁾.

Sample Preparation and Initial Cultivation: For each sample, 100 mL of drinking water were subjected to suction filtration through a filter paper (SKU: 09 790 2E). Once the suction process was complete, the filter paper was removed and brought into contact with specific media for the detection of pathogens: *Salmonella* spp. (XLD AGAR ISO 9001:2015 M4J2BV01 from TM Media), *Listeria* spp. (CHROMOGENIC LISTERIA AGAR BASE (Modified) M1GD4HV01 ISO 9001: 2015.ISO 11133:2014 from TM Media), and *E. coli* (EMB AGAR, LEVINE M3H1FV01 (ISO 21150:2015) from TM Media). The culture plates were incubated at a temperature of 35°C for 24 hours. After this time, colony counts and isolation were performed ⁽¹⁰⁾.

Confirmation by Microscopy: Plates that showed microbial growth were selected for Gram staining. By observing the morphology of the pathogen under a microscope, it was determined whether it was Gram-positive or Gram-negative ⁽¹¹⁾.

<https://orcid.org/0000-0001-9048-1538>

³ Grupo de investigación INVAGRO. Universidad Nacional de Chimborazo. Chimborazo. Ecuador.

<https://orcid.org/0000-0002-9066-6070>

⁴ Universidad Estatal de Bolívar, Facultad Ciencias Agropecuarias, Carrera Agroindustrias

<https://orcid.org/0000-0002-7434-4522>

DNA Extraction: Approximately 5 colonies of the pathogen were suspended in 500 μ L of 1X TAE buffer. The suspension was then centrifuged at 16,000 rpm for 10 minutes at room temperature. For DNA extraction of *Listeria* spp, *E. coli*, and both Gram-negative and Gram-positive *Salmonella* spp, the PurelinkTM Genomic DNA Mini Kit, K182001 ⁽¹²⁾, was used.

DNA Yield: To determine the DNA yield, a UV-Vis microvolume spectrophotometer, such as the Thermo ScientificTM NanoDropTM One/One, was utilized. It measures the concentration and purity of the samples, with measurements conducted on DNA concentrations from 1 to 1.5 μ L, and each sample's purity presented optimal percentages ⁽¹³⁾.

Preparation of *E. coli* Samples: A solution for *E. coli*, consisting of R Reverse (5'-GCTATTCCTGCCGATAAGAGA-3') with a concentration of 39.45 nmol and F Forward (5'-CCAGGCAAAGAGTTTATGTTGA-3') with a concentration of 37.39 nmol, was prepared as a working solution at a concentration of 1010 μ mol. Subsequently, for the preparation of the total sample solution, Nuclease-Free Water was added according to the nmol concentration. A 1.5 mL Eppendorf tube was used, and 5 μ L of the R primer were added, followed by topping up with 45 μ L of Nuclease-Free Water, resulting in a total solution of 50 μ L that recognizes the genus *Escherichia coli* with its serotype O157:H7 in 212 bp ⁽¹⁴⁾.

Sample Preparation for *Listeria monocytogenes*: The solution for *Listeria*, based on R Reverse (5'-GCCGTCGATGATTTGAACTTCATC-3') with a concentration of 41.88 nmol and F Forward (5'-GAATGTAAACTTCGGCGCAATCAG-3') with a concentration of 42.05 nmol, was prepared as a working solution at a concentration of 1010 μ mol. Nuclease-Free Water was added according to the nmol concentration. Using a 1.5 mL Eppendorf tube, 5 μ L of the R primer were added, and it was then filled up with 45 μ L of Nuclease-Free Water, resulting in a total solution of 50 μ L that recognizes the genus *Listeria monocytogenes* in 388 bp ⁽¹⁵⁾.

Sample Preparation for *Salmonella* spp. in Thermal Cycles: The solution for *Salmonella*, consisting of reverse invaA3R (5'-TCCATCAAATTAGCGGAGGC-3') with a concentration of 41.05 nmol and forwar inva3F (5'-AACGTGTTTCCGTCGTAAT-3') with a concentration of 52.15 nmol, was prepared as a final concentration of 1010 μ mol. Nuclease-Free Water was added. Using a 1.5 mL Eppendorf tube, 5 μ L of the R primer were added, and it was then filled up with 45 μ L of Nuclease-Free Water, resulting in a total solution of 50 μ L that recognizes the genus *Salmonella* spp. in 244 bp ⁽¹⁶⁾.

Table 1 Cebadores utilizados para realizar la PCR.

Bacteria	Sequence	Base pairs bp	References
Listeria monocytogenes	(R,5'-GCCGTCGATGATTTGAACTTCATC-3') (F,5'-GAATGTAAACTTCGGCGCAATCAG-3')	388 bp	(15)
Escherichia coli con su serotipo	(R,5'-GCTATTCCTGCCGATAAGAGA-3') (F,5'-	212 bp	(17)

O157:H7	CCAGGCAAAGAGTTTATGTTGA-3')		
Salmonella spp.	invaA3R (R,5'-TCCATCAAATTAGCGGAGGC-3') inva3F (F,5'-AACGTGTTTCCGTCGTAAT-3')	244 bp	(16)

Reagents for PCR

In the following tables, the quantities of reagents for use in PCR amplification of DNA obtained from three different microorganisms, namely *Listeria monocytogenes*, *Escherichia coli*, and *Salmonella* spp, are provided. The GoTaq® Green Master Mix, 2X, already contains a 2X solution ready for use. It includes DNA polymerase (GoTaq®), dNTPs, reaction buffer, MgCl, and blue and yellow colorants.

Table 2 PCR Reagents for the Studied Pathogens

Reactives	Volume uni.	<i>Listeria monocytogenes</i> # of Samples	<i>E. coli</i> # of Samples	<i>Salmonella</i> spp. # of Samples
GoTaq® Green Master Mix, 2X	12.5 µL			
Forwar primer, 10µM	0.25-2.5µL			
River, 10µM	0.25 2.5µL	20	40	10
DNA Templade	1-5 µL			
Nuclease-Free Wáter	25 µL			
to				
Total, Volume		830 /20=41,5 µL	1660 /40=41,5 µL	415 /10=41,5 µL

Note: Research Work.

Cycles for the Studied Pathogens in the Thermal Cycler

The cycles were conducted in a thermal cycler (Techne FTC3/05, series TC3000 20X0.5ML, origin Spain) as outlined in Table 3.

Table 3 DNA Amplification Cycles

N°of cycles	T (°C) / T <i>Listeria m.</i>	T (°C) / T <i>Salmonella spp.</i>	T (°C) / T <i>Escherichia coli</i>	Fases
1	95/2 min	95/5 min	95/30 s	Denaturation
	95/1 min	95/1 min	95/20 s	Denaturation
30	53/45 s	63/45 s	53/45 s	Primer Annealing
	72/ 1 min	72/1 min	72/1min	Extension
1	72/ 7 min	72/5 min	72/2min	Extension

Note: ⁽¹⁴⁾.

Upon completing these steps in the thermal cycler, a program is set up to carry out the specified cycles for the determined period.

Conventional PCR Electrophoresis:

A 1.5% Agarose gel was prepared in 100 mL of 1X TAE buffer. The gel was placed into the electrophoresis tank along with a 100 bp leader, and 5 µL of amplified DNA was added to each well. The electrophoresis run was conducted in the tank at 135 volts for 30 minutes. After the estimated time had passed, the gel was transferred to a staining tank, submerged in 1X TAE buffer with 1 µL of Diamond MT Nucleic Acid Dye for 15 minutes. The results were then read and interpreted using the Safe transilluminator photo documentation system, specifically the Invitrogen Serie 12078086 transilluminator, with the aid of a UV transilluminator for visualization ⁽¹⁸⁾.

RESULTS:

Georeferencing of Different Sampling Points in Chambo Canton:

Table number 4 presents the georeferencing results achieved using a GPS MAP 64sx through ArcGIS software. This process allowed for the precise location of sampling points within Chambo Canton by using specific coordinates.



Figure 1: Sampling Points in the Urban Area of Chambo Canton.

Table 4 Sampling Points for Human Consumption Water in Chambo Canton.

Sampling Points	Geographic Area
P1, P2, P3, P4, P5	Northwest - Mountainous Area - Sawmill Zone and Storage Tank
P6, P7, P8, P9, P10, P11	Northwest - Mountainous Area - Springs - Storage Tank - 200 meters from Cubillin Estate
P12, P13, P14, P16	Northwest - Storage Tanks - Treatment Tanks - Distribution Tanks
P17, P18, P19, P20, P21, P22, P23, P24	Southwest - First Houses with Potable Water
P25, P26, P27, P28, P29, P30, P32, P33, P34, P50	Southwest - Manuel Zabala and Egidio Fierro Streets
P35, P36, P37, P38, P39	Southeast - Héctor Guevara and 18 de Marzo Streets
P40, P41, P42, P43, P44	Southwest - Héctor Guevara - Edelberto Bonilla - David Parra

P45, P46, P47, P49

Southwest - F Street - Chiriboga -
Rocafuerte and García Moreno Streets

Table 4 displays the water sampling points in Chambo Canton, which were collected from areas in the northeastern part of the city, including springs, streams, and rivers. Additionally, in the southwestern and southeastern areas, various points were sampled where treated water is distributed for human consumption.

The microbiological results of the samples from the different sampling points in Chambo Canton

Table 5 Microbiological Results.

Code	monocytogenes	Salmonella	Reference	Analysis	Method
Listeria	E. Coli	Unit			
P1	-	+	+		
P2	+	+	+		
P3	+	+	+		
P4	-	+	-		
P5	-	+	-		
P6	+	+	+		
P7	+	+	+		
P8	+	+	+	CFUs./	ISO
P9	+	+	+	100mL	9001:2015
P10	+	+	+		
P11	+	+	+		
P12	+	+	+		
P13	+	+	-		
P14	-	-	-		
P15	-	-	-		
P16	-	-	-		
P17	-	-	-		
P18	-	-	-		
P19	+	-	-		
P20	+	-	-		
P21	-	-	-		
P22	-	-	-		
P23	+	+	-		
P24	+	+	-		
P25	-	+	-		
P26	-	+	-		
P27	-	+	-		
P28	+	+	-		
P29	-	+	-		

P30	-	+	-
P31	+	+	-
P32	-	+	-
P33	-	+	-
P34	-	+	-
P35	+	+	-
P36	-	+	-
P37	-	+	-
P38	-	+	-
P39	-	+	-
P40	+	+	-
P41	-	+	-
P42	-	+	-
P43	-	+	-
P44	+	+	-
P45	-	+	-
P46	+	+	-
P47	-	+	-
P48	-	+	-
P49	-	+	-
P50	-	+	-

Table 5 presents the results of the sampling of drinking water from different points in Chambo Canton. These results were obtained through microbiological analyses using specific and selective media. It's worth noting that the analyses were conducted in triplicate during the research process to confirm their prevalence.

Table 6 It appears to deal with the microscopic identification and differentiation, likely based on Gram staining, following the microbiological results of the drinking water samples.

Code	Listeria monocytogenes	E. Coli	Salmonella
P1	-	Bacilos (-)	Bacilos (-)
P2	Bacilos (+)	Bacilos (-)	Bacilos (-)
P3	Bacilos (+)	Bacilos (-)	Bacilos (-)
P4	-	Bacilos (-)	-
P5	-	Bacilos (-)	-
P6	Bacilos (+)	Bacilos (-)	Bacilos (-)

P7	Bacilos (+)	Bacilos (-)	Bacilos (-)
P8	Bacilos (+)	Bacilos (-)	Bacilos (-)
P9	Bacilos (+)	Bacilos (-)	Bacilos (-)
P10	Bacilos (+)	Bacilos (-)	Bacilos (-)
P11	Bacilos (+)	Bacilos (-)	Bacilos (-)
P12	Bacilos (+)	Bacilos (-)	Bacilos (-)
P13	Bacilos (+)	Bacilos (-)	-
P14	-	-	-
P15	-	-	-
P16	-	-	-
P17	-	-	-
P18	-	-	-
P19	Bacilos (+)	-	-
P20	Bacilos (+)	-	-
P21	-	-	-
P22	-	-	-
P23	Bacilos (+)	-	-
P24	Bacilos (+)	Bacilos (-)	-
P25	-	Bacilos (-)	-
P26	-	Bacilos (-)	-
P27	-	Bacilos (-)	-
P28	Bacilos (+)	Bacilos (-)	-
P29	-	Bacilos (-)	-
P30	-	Bacilos (-)	-
P31	Bacilos (+)	Bacilos (-)	-
P32	-	Bacilos (-)	-
P33	-	Bacilos (-)	-
P34	-	Bacilos (-)	-
P35	Bacilos (+)	Bacilos (-)	-

P36	-	Bacilos (-)	-
P37	-	Bacilos (-)	-
P38	-	Bacilos (-)	-
P39	-	Bacilos (-)	-
P40	Bacilos (+)	Bacilos (-)	-
P41	-	Bacilos (-)	-
P42	-	Bacilos (-)	-
P43	-	Bacilos (-)	-
P44	Bacilos (+)	Bacilos (-)	-
P45	-	Bacilos (-)	-
P46	Bacilos (+)	Bacilos (-)	-
P47	-	Bacilos (-)	-
P48	-	Bacilos (-)	-
P49	-	Bacilos (-)	-
P50	-	Bacilos (-)	-

Table 7 The concentration and quality of extracted DNA from microorganisms like *Listeria monocytogenes*, *E. coli*, and *Salmonella* spp.

Code	<i>Listeria monocytogenes</i> ng/ μ L	<i>E. Coli</i> ng/ μ L	<i>Salmonella</i> ng/ μ L
P1	-	56,8	69,5
P2	67,5	70,5	51,6
P3	98,2	68,9	87,4
P4	-	93,4	-
P5	-	76,7	-
P6	73,9	87,6	83,7
P7	68,3	101,8	78,9
P8	80,5	99,4	56,8
P9	48,9	102,5	64,8
P10	56,4	89,1	89,7
P11	78,5	97,2	47,9

P12	98,2	83,9	55,6
P13	87,4	68,7	-
P14	-	-	-
P15	-	-	-
P16	-	-	-
P17	-	-	-
P18	-	-	-
P19	102,3	-	-
P20	55,1	-	-
P21	-	-	-
P22	-	-	-
P23	87,3	54,7	-
P24	49,2	77,2	-
P25	-	94,6	-
P26	-	48,9	-
P27	-	62,4	-
P28	78,3	70,6	-
P29	-	58,9	-
P30	-	73,5	-
P31	67,4	54,6	-
P32	-	107,2	-
P33	-	65,4	-
P34	-	48,2	-
P35	63,5	75,8	-
P36	-	76,1	-
P37	-	64,9	-
P38	-	79,1	-
P39	-	89,7	-
P40	54,6	92,4	-
P41	-	100,2	-
P42	-	59,7	-
P43	-	74,8	-
P44	76,8	34,5	-
P45	-	76,0	-
P46	34,5	78,9	-
P47	-	56,7	-
P48	-	49,8	-
P49	-	56,7	-
P50	-	52,7	-

Results of PCR Amplification via Agarose Gel Electrophoresis for *Escherichia coli*, *Listeria monocytogenes*, and *Salmonella*.

Results obtained from photographic documentation on an agarose gel with isolates of

Escherichia coli and its serotype O157:H7, considering a positive control (E+), with various points (P) containing amplified DNA, in comparison to a 100 bp molecular weight marker.




Figure 2: Represents the contaminated samples from P1 to P28.




Figure 3: Represents the contaminated samples from P29 to P46.



Figure 4: Represents the contaminated samples from P46 to P50




Figure 5: Results obtained from photographic documentation on an agarose gel with isolates of *Listeria monocytogenes* and its serotype O157:H7, considering a positive control (E+), with various points (P) containing amplified DNA, in comparison to a 100 bp molecular weight marker, represents the contaminated samples from P2 to P40.




Figure 5: Represents the contaminated samples from P44 to P46.




Figure 7: Results obtained from the photographic documentation on an agarose gel with isolates of *Salmonella* spp. and its serotype O157:H7, considering a positive control (E+), with various points (P) containing amplified DNA, compared to a 100 bp molecular weight marker, represents the contaminated samples from P1 to P12.

DISCUSSION

Water is a fundamental element for human biological activities, and its quality must be suitable for consumption to prevent diseases that can harm consumers and even lead to fatalities ⁽¹⁹⁾. In this research, the quality of drinking water in Chambo Canton was evaluated, focusing on the microbiology of the water matrix based on the standard ⁽²⁰⁾. Based on the data obtained, it is concluded that this vital liquid is not suitable for consumption.

Pathogens that can cause foodborne illnesses with gastrointestinal symptoms include *Listeria monocytogenes*, *E. Coli*, and *Salmonella* spp. ⁽²¹⁾. Among the 50 points studied, it was observed that the presence of *E. Coli* was detected in 40 samples, representing 80%, *Listeria monocytogenes* was found in 20 out of 50 samples, accounting for 40%, and *Salmonella* spp. was present in 10 samples out of 50, constituting 10%. This suggests that the studied water may potentially transmit diseases such as typhoid fever, paratyphoid fever, and hepatitis A ⁽²²⁾.

CONCLUSIONS

Through georeferencing, the position of water samples was analyzed using GPS coordinates and processed with ArcGIS software. Based on the georeferenced points, it can be concluded that the journey of human consumption water is contaminated both before and after treatment.

There is contamination with *Escherichia coli* of serotype O157:H7 in 80% of the samples (40 out of 50), followed by *Listeria monocytogenes* at 40% (20 out of 50), and *Salmonella* spp. at a rate that isn't specified in the provided text. These findings indicate a significant risk to public health and highlight the need for immediate action to address water contamination in Chambo Canton.

20% (10/50), thus providing a reason to acknowledge that the drinking water in Chambo is contaminated by the studied bacteria. It can be considered that the water, by not complying with the INEN 1108 and INEN 1107 standards, may be influenced by factors such as the deterioration of the drinking water distribution network and the proximity of networks for other uses and wastewater to this network.

Knowing that there is contamination from pathogens that are not heat-resistant, it is recommended to treat drinking water with thermal processing, boiling it at temperatures ranging from 98°C to 100°C. Alternatively, a chemical purification method using

chlorine at a minimum concentration of 0.2 mg/L and a maximum of 2 mg/L, as per INEN 409-03: 2021, can be employed.

It is important that local and regional authorities take immediate action to address this contamination and ensure access to safe drinking water for the population of Chambo. Additionally, ongoing water quality monitoring and preventive measures should be implemented to prevent future contamination.

CONFLICT OF INTERESTS

The authors who conducted this research do not have any conflicts of interest.

References

1. Lopez Ce. El agua para una transformación social-ecológica. FES Transformacion. 2018;; p. 79.
2. ACUAE F. ACUAE FOUNDATION. [Online].; 2021. Acceso 30 de 09 de 2023. Disponible en: <https://www.fundacionaquae.org/wiki/cantidad-de-agua-potable-fuente-de-vida/#:~:text=%C2%BFCu%C3%A1%20es%20la%20cantidad%20de,la%20Tierra%20se%20considera%20dulce.>
3. UNESCO. Greenfact. [Online].; 2019. Acceso 30 de 09 de 2023. Disponible en: www.unesco.org/water/wwap/wwdr2/pdf/wwdr2_ch_4.pdf.
4. OPS oPDS. paho.org. [Online].; 2022. Acceso 30 de 09 de 2023. Disponible en: <https://www.paho.org/es/temas/agua-saneamiento>.
5. Shayo GM. Severity of waterborne diseases in developing countries and the effectiveness of ceramic filters for improving water quality. Bulletin of the National Research Centre. 2023; 47(113). <https://doi.org/10.1186/s42269-023-01088-9>
6. Barrera G. PCR en tiempo real: una metodología útil para la detección y cuantificación de granulovirus. Rev. Colomb. Biotecno. 2016; 18(2). <https://doi.org/10.15446/rev.colomb.biote.v18n2.61514>
7. Mejía LM. Análisis microbiológico del agua para consumo humano de la población del centro poblado pachapiriana, distrito de chontalí. Ciencia Latina Revista Científica Multidisciplinar. 2021; 5(6). https://doi.org/10.37811/cl_rcm.v5i6.1355
8. Alvarez Y. Georreferenciación de documentos cartográficos históricos para el análisis del trazado fluvial del bajo segura, vega media (MURCIA, ESPAÑA). 2018;(21): p. 101-118. <https://doi.org/10.21138/GF.536>
9. INEN2176 N. Agua. Calidad del agua muestreo. tecnicas de muestreo..
10. Sánchez EP. Simulación y Conteo de Unidades Formadoras de colonias. Computacion e informatica. 2017; 6(1): p. 97-111.
11. Corrales LC. Principios físicoquímicos de los colorantes utilizados en microbiología. NOVA. 2020; 18(33): p. 73-100. <https://doi.org/10.22490/24629448.3701>
12. Villota GE. Aislamiento y caracterización de bacterias productoras de biopolímeros a partir de efluentes industriales. Colomb. Biotecnol. 2022; 14(1): p. 27-45. <https://doi.org/10.15446/rev.colomb.biote.v24n1.76660>
13. Mancera P. Cuantificación de ácidos nucleicos mediante diferentes técnicas..
14. Yepes M. Validación de PCR convencional para detectar Escherichia coli O157. Información Tecnológica. 2022; 33(2): p. 3-12. <https://doi.org/10.4067/S0718-07642022000200003>
15. Meghdadi H, Nassirabady N. Isolation and characterization of Listeria monocytogenes from. Bogota: Ahvaz Jundishapur University of Medical Sciences, Facultad de Microbiología.
16. Gomez J, Torres J. Prevalencia y diversidad genómica de Salmonella enterica recuperada

del agua de un río en una importante región agrícola en el noroeste de México. Culiacan-Mexico: Centro de Investigación en Alimentación y Desarrollo A.C. (CIAD), Laboratorio Nacional para la Investigación en Inocuidad Alimentaria.

17. Nuñez D, Bayas F. Prevalencia de Salmonella Spp. y Escherichia Coli en agua de consumo humano en Ecuador. Guaranda-Chimbo: Universidad Estatal de Bolivar.
18. Wahjudi M. Detection of Salmonella spp., Escherichia coli, and Listeria monocytogenes in Tuna by Multiplex PCR. Journal of Aquaculture and Fish Health. 2022; 11(3): p. 391-401. <https://doi.org/10.20473/jafh.v11i3.36509>
19. Guzmán BL. La calidad del agua para consumo humano y su asociación con la morbimortalidad en Colombia, 2008-2012. Biomédica. 2015; 35(2): p. 177-190. <https://doi.org/10.7705/biomedica.v35i0.2511>
20. INEN1108. AGUA POTABLE. REQUISITOS..
21. Torres YF. Patógenos asociados a enfermedades transmitidas por alimentos en restaurantes escolares de Colombia. Rev Chil Nutr. 2017; 44(4). <https://doi.org/10.4067/S0717-75182017000400325>
22. Miranda JPR. Enfermedades transmitidas por el agua y saneamiento básico en Colombia. Rev. Salud Pública. 2016; 18(5): p. 738-745. <https://doi.org/10.15446/rsap.v18n5.54869>
23. Darshini M, Pope M. PureLink DNA (invitrogen by life technologies USA; 2012.
24. OMS. [Online].; 2023. Acceso 30 de 09 de 2023. Disponible en: <https://www.who.int/es/news-room/fact-sheets/detail/drinking-water#:~:text=E1%20agua%20para%20consumo%20humano,000%20muertes%20por%20enfermedades%20diarreicas.>
25. NTE INEN 2176 Pr. [Online].; 2013. Acceso 30 de 09 de 2023. Disponible en: https://gestionambiental.pastaza.gob.ec/biblioteca/legislacion-ambiental/patrimonio_natural/nte_inen_2176_1_agua_calidad_agua_muestreo_tecnicas_muestreo.pdf.
26. ISO 7218 U. saludneuquen. [Online].; 2008. Acceso 30 de 09 de 2023. Disponible en: https://www.saludneuquen.gob.ar/wp-content/uploads/2021/08/UNE-EN_ISO_72182008.pdf.