

Bioremediation Strategies For Soil And Water Pollution Harnessing The Power Of Microorganisms

Dr. Saurabh Sanjay Joshi¹, Mr. Pramod Keshavrao Jadhav², Dr. Naveen K Chourasia³, Mr. Prasad Janardhan Jadhav⁴, Dr. Khushal N. Pathade⁵, KANIMOZHI K R⁶

Abstract:

Environmental problems such as soil and water contamination are common and are caused by human activities such as industrial processes and agricultural practices. These pollutants include a wide range of chemical contaminants, such as pesticides, heavy metals, pesticide-containing chlorinated solvents, polycyclic aromatic hydrocarbons (PAHs), and newly developing contaminants like microplastics and medicines. Urgent remedial action is required due to the detrimental effects of such pollutants on human health, ecological integrity, and socio-economic welfare. Utilising the innate metabolic capacities of microorganisms to convert and detoxify pollutants into safe metabolites, bioremediation has become a compelling and long-lasting method for mitigating pollution. The present study offers a thorough examination of bioremediation tactics designed to tackle soil and water contamination, focusing on the complex interactions among microbial communities, environmental factors, and remediation effectiveness.

Introduction

Microbial consortia, which are composed of bacteria, fungus, archaea, and algae, are the fundamental components of bioremediation. These organisms possess distinct enzymatic repertoires that enable them to catalyse a wide range of bio¹transformation events. These microorganisms metabolise resistant contaminants and assimilate them into harmless end products or cellular biomass by using a variety of metabolic pathways, such as co-metabolism, fermentation, aerobic and anaerobic respiration, and enzymatic destruction. Numerous bioremediation strategies are explained, including in situ and ex situ methods suited to particular contaminant matrices and environmental circumstances. Exogenous microbial inocula are introduced in bioaugmentation strategies to increase substrate specificity or speed up degradation, while environmental factors like redox potential and nutrient availability are adjusted in biostimulation strategies to increase native microbial activity. Microbial fuel cells

¹Head and Associate Professor, Department of Civil and Environmental Engineering, Kolhapur Institute of Technology's College of Engineering (Autonomous), Kolhapur

²Assistant Professor, Department of Civil and Environmental Engineering, Kolhapur Institute of Technology's College of Engineering (Autonomous), Kolhapur.

³Assistant Professor, Dept of Zoology, Government College, Bichhua, Chhindwara

⁴Assistant Professor, Department of Civil and Environmental Engineering, Kolhapur Institute of Technology's College of Engineering (Autonomous), Kolhapur.

⁵Assistant Professor and Head P.G. Department of Botany Dr. R. G. Bhojar Arts, Commerce and Science College, Seloo Dist. Wardha, Maharashtra

⁶Associate Professor Department of Chemistry Sri Krishna College of Technology Coimbatore 641042

use microbial electrochemical processes to produce energy and remove pollutants, while phytoremediation uses the phytotransformation and rhizosphere-mediated mechanisms of plants to accelerate the uptake, translocation, and breakdown of pollutants.

Numerous variables, such as the physicochemical characteristics of the pollutants (such as solubility, volatility, and bioavailability), the environmental factors (such as pH, temperature, and moisture content), the makeup of the microbial community, and interactions with co-contaminants, can affect how effective bioremediation is. Our understanding of microbial ecology and metabolic networks has been completely transformed by developments in molecular microbiology, omics technologies, and bioinformatics. This has made it possible to manipulate and optimise bioremediation processes in a targeted manner.

Bioremediation holds great potential, but it faces many obstacles along the way, such as substrate inhibition, microbial competition, mass transfer restrictions, and the rise of microbial pathogens and antibiotic-resistant genes. In addition, the enduring nature of resistant contaminants and the intricate relationships among microbial groups need the use of integrated strategies that incorporate physical, chemical, and thermal treatments with bioremediation.

The analysis concludes by highlighting the critical role that bioremediation plays in tackling soil and water contamination, providing a sustainable and environmentally acceptable substitute for traditional remediation methods. Through the utilisation of microbial communities' metabolic capabilities, bioremediation presents significant opportunities for the restoration of damaged environments and the preservation of ecosystem health. Sufficient multidisciplinary research, technological advancements, and governmental backing are essential for actualizing the complete potential of bioremediation and reducing the widespread effects of soil and water contamination on the sustainability of the global environment.

The tremendous diversity of microorganisms, including bacteria, fungus, algae, archaea, and protozoa, which are each endowed with distinct metabolic pathways and enzyme repertoires suited to particular classes of contaminants, is the fundamental component of bioremediation. These bacteria may metabolise a broad variety of organic and inorganic pollutants because to their extensive metabolic capacities, which include co-metabolism, enzymatic degradation, aerobic and anaerobic respiration, and fermentation. Moreover, metabolic byproducts from one species can act as substrates or co-factors for other species in microbial consortia, improving the overall effectiveness of remediation. This phenomenon is known as synergistic interaction.

Many bioremediation strategies are explained; these include ex situ procedures, which remove and treat contaminated matrices in a controlled environment, and in situ procedures, which treat contaminants in their natural habitat. Biostimulation techniques alter environmental parameters (e.g., redox conditions, nutrient availability) to stimulate indigenous microbial activity, whereas bioaugmentation strategies introduce exogenous microbial inocula to increase degradation rates or broaden substrate specificity. Through biochemical processes like phytotransformation, rhizodegradation, and phytostabilization, plants have the unique ability to absorb, translocate, and metabolise pollutants. This process is known as phytoremediation. Furthermore, microbial fuel cells provide a dual-purpose method for energy production and environmental remediation by using microbial electrochemical activities to produce power and degrade organic contaminants at the same time.

This analysis concludes by highlighting the vital significance of bioremediation as a long-term, environmentally responsible solution to soil and water contamination. Through the utilisation of microbial communities' metabolic capacities, bioremediation provides economical and

proficient methods for cleaning contaminated areas, all the while reducing ecological disturbance and enhancing the resilience of ecosystems. To fully use bioremediation and lessen the widespread effects of soil and water pollution on the sustainability of the environment worldwide, multidisciplinary research, technological advancement, and policy assistance are crucial

1. Introduction

Because of the unrelenting advances in industry, urbanisation, intensification of agriculture, and human activities, soil and water contamination have alarmingly increased in the 21st century (1). The range of contaminants that have proliferated includes heavy metals, polycyclic aromatic hydrocarbons (PAHs), pesticides, herbicides, herbicides, pharmaceuticals, endocrine disruptors, and nanoplastics. These pollutants have impacted both terrestrial and aquatic ecosystems, causing damage to soils, sediments, surface waters, and aquifers.

Pollution of soil and water has numerous negative effects on human health, the environment, and socioeconomic well-being (2). Contamination causes long-lasting disturbances to food web dynamics, nutrient cycling, and ecological succession, which in turn cause loss of biodiversity, habitat degradation, and dysfunction in ecosystems. Contaminants bioaccumulate and biomagnify through trophic levels, posing dangers to wildlife and human populations, with severe health impacts range from acute toxicity and mutagenesis to cancer and reproductive abnormalities (3). Moreover, pollution-induced perturbations in ecosystem services, including water purification, soil fertility, temperature control, and carbon sequestration, produce cascading repercussions on global biogeochemical cycles and planetary health.

Bioremediation has become a persuasive and environmentally sound method of mitigating soil and water contamination in response to the pressing need for remedial action (4). By utilising the inherent biochemical capacities of microbial populations to break down, modify, and remove pollutants, bioremediation makes use of the metabolic powers of microorganisms and provides a viable and affordable substitute for traditional remediation techniques.

The complex metabolic networks and enzymatic repertoires of microorganisms—which include bacteria, fungus, algae, archaea, and protozoa—that are each equipped with unique mechanisms for pollutant degradation and assimilation are essential to the effectiveness of bioremediation (5). These microorganisms metabolise a wide range of organic and inorganic pollutants, converting them into harmless end products or assimilating them into cellular biomass (6). They do this by using a variety of metabolic strategies, such as co-metabolism, enzymatic degradation, fermentation, and aerobic and anaerobic respiration (7).

A number of factors come together to determine whether bioremediation is successful. These include the physicochemical characteristics of the contaminants (such as hydrophobicity, solubility, and persistence), the environmental factors (such as pH, temperature, and redox potential), the makeup of the microbial community, the availability of substrate, and interactions with co-contaminants. Furthermore, the particular contamination matrices and environmental contexts influence the choice and optimisation of bioremediation strategies, such as phytoremediation, rhizoremediation, mycoremediation, bioaugmentation, and biostimulation (8).

This work strives to present a thorough overview of bioremediation solutions for soil and water contamination, with a special emphasis on the crucial role of microorganisms in driving remediation processes (9). In order to understand the complexities of microbial-mediated biotransformation processes, it will draw on recent developments in molecular microbiology,

genomics, metagenomics, transcriptomics, proteomics, metabolomics, and bioinformatics (10). This will clarify the underlying biochemical mechanisms and metabolic pathways used by microbial consortia to degrade various classes of pollutants (11).

This paper aims to demonstrate the practical applicability and efficacy of bioremediation in real-world circumstances by analysing case studies and field trials that showcase successful bioremediation initiatives in various environmental conditions. This review aims to inform and inspire future research, innovation, and policy initiatives aimed at fostering environmental sustainability and resilience in the face of increasing pollution pressures by clarifying the basic concepts, technological advancements, and ecological implications of bioremediation.

As a powerful and promising solution to reduce soil and water pollution, bioremediation offers a paradigm shift towards ecological resilience and sustainable environmental stewardship (12). We can unlock the revolutionary potential of bioremediation and effect the rehabilitation and restoration of damaged habitats for the benefit of current and future generations by utilising the metabolic inventiveness of microorganisms.

Methodology

2.1 Literature Review:

The research paper's methodology comprises a thorough and methodical assessment of the literature to compile the state of the art and new developments in bioremediation techniques for reducing pollution in the soil and water (13). A thorough search of electronic databases, including PubMed, Web of Science, Scopus, Google Scholar, and specialised environmental science databases, was carried out using a multidisciplinary approach. To guarantee the retrieval of pertinent literature, a broad range of keywords and controlled vocabulary terms were used, such as "bioremediation," "microbial degradation," "phytoremediation," "bioaugmentation," "biostimulation," "microbial fuel cells," "contaminant removal," "pollutant fate," and "environmental microbiology."

2.2 Data Collection and Synthesis:

The process of gathering data entailed a methodical retrieval of academic sources pertaining to environmental microbiology and bioremediation, including books, reports, conference proceedings, and peer-reviewed articles. Relevant data on bioremediation methods, microbiological processes, pollutant kinds, environmental conditions, case studies, and technological advancements were retrieved from a selection of literature sources that were rigorously evaluated. In order to identify patterns, trends, and gaps in the literature, data were sorted, classified, and synthesised. Where appropriate, qualitative and quantitative analytical techniques were used.

2.3 Analysis and Interpretation:

Using a variety of analytical methods, such as content analysis, theme analysis, comparative analysis, and statistical analysis, the gathered data was thoroughly examined (14). While quantitative analysis included the tabulation, classification, and statistical processing of numerical data such as pollutant removal efficiencies, microbial diversity indices, and environmental parameters, qualitative analysis focused on identifying recurrent themes, concepts, and theoretical frameworks found in the literature. The findings were analysed in

light of current theories, models, and empirical data in order to produce insightful deductions and conclusions that are supported by science (15).

2.4 Integration of Findings:

The results of the literature review were combined and refined to create a cohesive story that summarises the most recent advancements in bioremediation science (16). Clarifying the fundamental mechanisms of microbial-mediated pollutant degradation, assessing the limitations and effectiveness of various bioremediation techniques, determining the factors influencing bioremediation performance, and identifying new directions and research gaps in the field were all priorities. The process of synthesising findings made it easier to uncover similarities and differences amongst studies, which allowed for a thorough grasp of the topic and informed the creation of research recommendations and policy implications (17).

2.5 Gap Analysis and Future Directions:

To determine knowledge gaps, research objectives, and opportunities for further study in the field of bioremediation, a critical gap analysis was carried out. Finding gaps in the body of literature provided the foundation for suggestions for new lines of inquiry, developments in technology, and interdisciplinary partnerships with the goal of promoting bioremediation as a science and practice. Based on the synthesis of data from the literature study, recommendations were made for future research areas that would address identified gaps, optimise bioremediation technologies, enhance remediation efficiency, and promote sustainable environmental management practices.

2.6 Ethical Considerations:

Ensuring ethical issues were of utmost importance during the research process involved adhering to standards of academic integrity, transparency, and responsibility. In order to recognise the contributions of earlier research and scholarship, proper citation and attribution of sources were made sure to do. A balanced and objective examination of the literature was presented, without prejudice or undue influence from external parties. Potential conflicts of interest were declared. In all facets of data collecting, processing, and reporting, confidentiality, privacy, and intellectual property rights were respected.

2.7 Quality Assurance and Peer Review:

To maintain academic rigour, credibility, and dependability, the methodology and findings of this research study were subjected to stringent quality assurance and peer review procedures (18). The research technique and the robustness of the findings were validated by subjecting the methodology to rigorous scrutiny from mentors, academic colleagues, and specialists in environmental science and bioremediation. Peer reviewers' comments was requested and taken into consideration to improve the research paper's clarity, coherence, and comprehensiveness as well as to reinforce the research technique.

1. Understanding Soil and Water Pollution

The introduction of pollutants into terrestrial and aquatic ecosystems gives birth to complex and multidimensional environmental concerns, such as soil and water contamination (19). Numerous human activities, including as industrial processes, agriculture, urbanisation, mining, transportation, and waste disposal, are the source of these pollutants (20). These activities lead to the infiltration of a wide range of chemical, physical, and biological

contaminants into soil and water ecosystems, which poses serious concerns to biodiversity, ecosystem health, and human welfare.

3.1 Sources and Types of Pollutants:

Pollution of soil and water comes from a variety of point and non-point sources. Point sources are locations where pollutants are released into the environment directly, such as chemical spills, hazardous waste sites, wastewater treatment facilities, and industrial discharge pipes. Conversely, diffuse inputs—such as urban storm water runoff, agricultural runoff, atmospheric deposition, and groundwater contamination—are examples of non-point sources since they include the spread of contaminants over wide regions via a variety of routes.

Both soil and water can include a wide variety of pollutants, from organic chemicals and heavy metals to microbial diseases and newly discovered toxins (21). Common industrial activities including mining, smelting, metal plating, and waste incineration can release heavy metals like lead, cadmium, mercury, arsenic, chromium, and nickel (22). Due to their bioaccumulation and biomagnification, these metals are hazardous to ecosystems and human health because they are persistent in the environment and can build up in soil and sediment (23).

Polycyclic aromatic hydrocarbons (PAHs), chlorinated solvents, insecticides, herbicides, industrial chemicals, medicines, and personal care items are just a few examples of the many synthetic chemicals that fall under the category of organic pollutants. These substances can remain in the environment for extended periods of time and pose a risk to aquatic life, wildlife, and human health. They are obtained from industrial processes, agricultural methods, household goods, and pharmaceuticals.

Faecal pollution, agricultural runoff, sewage discharges, and animal waste are some of the ways that microbial pollutants—which include bacteria, viruses, protozoa, and helminths—can pollute soil and water. These pathogens increase the risk of contracting waterborne illnesses like viral gastroenteritis, cholera, typhoid fever, dysentery, giardiasis, and cryptosporidiosis, especially in places with poor water quality management and inadequate sanitary facilities.

A novel class of pollutants known as emerging contaminants has drawn attention because of the possible harm they could do to the environment and public health (24). Among these pollutants include flame retardants, medicines, endocrine-disrupting chemicals (EDCs), nanomaterials, microplastics, and per- and polyfluoroalkyl substances (PFAS). Novel techniques for detection, evaluation, and remediation may be necessary when emerging pollutants evade conventional monitoring and treatment techniques.

3.2 Environmental Fate and Transport:

A complex interplay of physical, chemical, biological, and hydrological processes controls the fate and transit of contaminants in soil and water habitats. Many pollutants find their primary sink in soil, where their physicochemical qualities and environmental conditions can cause them to volatilize into the atmosphere, attach to soil particles, or become sequestered in soil organic matter or leak into groundwater.

Contaminants in soil can go through a variety of processes that affect their mobility, bioavailability, persistence, and transformation, including adsorption, desorption, volatilization, dissolution, precipitation, sorption, leaching, and biodegradation. Important roles are played by soil characteristics in modulating pollutant behaviour and fate, including texture, structure, composition, pH, moisture content, redox potential, and microbial activity

(25).

Pollutants can disperse, dilute, advection, sedimentation, and bioaccumulate in aquatic habitats, impacting aquatic ecosystems and posing health concerns to aquatic creatures and humans (26). Surface water runoff, groundwater movement, atmospheric deposition, and aquatic currents are some of the ways that pollutants can be carried and end up contaminating surface waters, groundwater, sediments, and aquatic biota (27).

Pollutant routes and fates in aquatic ecosystems are influenced by hydrological processes such as runoff, infiltration, percolation, evaporation, transpiration, and surface water flow dynamics (28). Pollutants can have long-term ecological effects and pose dangers to human health via bioaccumulating in sediments, bioaccumulating in aquatic creatures, and biomagnifying in food webs.

3.3 Ecological and Human Health Impacts:

Pollution of soil and water affects ecosystems, biodiversity, human health, and socioeconomic well-being at several regional and temporal scales. Disruptions to trophic interactions, energy flow, nutrient cycling, species composition, and ecosystem services can result in ecosystem services, biodiversity, and ecological resilience being reduced (29). Through food webs, contaminants have the ability to bioaccumulate and biomagnify, endangering human populations as well as wildlife through direct contact or ingestion of polluted food and water supplies. Numerous detrimental health impacts, such as respiratory conditions, neurological diseases, developmental abnormalities, reproductive impairments, immunological disorders, and carcinogenicity, can arise from long-term exposure to pollution (30).

Additionally, soil and water pollution can worsen social injustices by disproportionately harming marginalised groups, low-income communities, indigenous peoples, and other vulnerable populations that lack access to clean water, sanitary facilities, and medical treatment. The financial consequences of health care expenses linked to pollution, environmental cleanup costs, lost productivity, and reduced ecosystem services highlight how urgent it is to address soil and water pollution using comprehensive and integrated strategies.

In conclusion, soil and water pollution provide serious obstacles to human health, socio-economic growth, and environmental sustainability (31). As a result, coordinated efforts are required to reduce pollution sources, clean up polluted areas, and advance sustainable resource management techniques. We can create well-informed strategies for pollution prevention, regulation, monitoring, and remediation, protecting environmental quality and guaranteeing the welfare of current and future generations, by knowing the origins, kinds, fate, transport, and impacts of pollutants in soil and water.

4. Importance of Bioremediation

Bioremediation is a very important environmentally responsible method of treating soil and water contamination. Several significant elements highlight its importance:

- 4.1 Environmental Sustainability:** Bioremediation offers a sustainable alternative to traditional remediation methods including excavation, incineration, and chemical treatment, which usually have a significant detrimental impact on the environment and result in high energy and carbon emissions. Using natural biological processes,

bioremediation minimises ecological disturbance, preserves soil fertility, safeguards biodiversity, and increases ecosystem resilience. It is consistent with the ideas of environmental sustainability and conservation.

- 4.2 Cost-Effectiveness:** Because bioremediation doesn't require costly infrastructure, equipment, or chemical inputs, it is frequently less expensive than conventional remediation procedures. As the fundamental agents of bioremediation, microorganisms are widely distributed, self-replicating, and, once established in contaminated places, require little in the way of maintenance or operating expenses. Remediation costs can be further decreased by using bioremediation in situ, which eliminates the requirement for site excavation, transportation, and disposal of polluted soil or water.
- 4.3 Versatility and Applicability:** Among the many toxins that can be removed from soil and water via bioremediation include pesticides, heavy metals, organic pollutants, petroleum hydrocarbons, and newly discovered chemicals. Numerous bioremediation techniques, including phytoremediation, bioaugmentation, biostimulation, and microbial fuel cells, provide customised answers for various pollutant kinds, site-specific difficulties, and environmental circumstances. Because of its adaptability, bioremediation can be used in a variety of environmental contexts, including urban landscapes, industrial brownfields, agricultural fields, and aquatic environments.
- 4.4 Minimization of Secondary Environmental Risks:** In contrast to certain chemical-based remediation techniques, which could produce toxic byproducts or increase secondary environmental concerns, bioremediation provides a safe and non-invasive way to reduce pollution. Pollutants are broken down by microorganisms into less hazardous, simpler molecules that are frequently integrated into their biomass or mineralized into innocuous byproducts like carbon dioxide, water, and biomass. This lessens the possibility of long-term environmental liabilities by lowering the likelihood of pollutant leaching, volatilization, or mobilisation into nearby soil, water, or air compartments.
- 4.5 Compatibility with Sustainable Development Goals:** Since bioremediation addresses several facets of environmental, social, and economic sustainability, it is consistent with the goals of sustainable development. Through the promotion of clean water resources, the restoration of terrestrial ecosystems, and the conservation of biodiversity, it helps to achieve Goal 6 (Clean Water and Sanitation) and Goal 15 (Life on Land) of the United Nations Sustainable Development Goals (SDGs). Furthermore, via encouraging technological innovation and developing sustainable practices in environmental management and remediation, bioremediation contributes to the achievement of Goal 9 (Industry, Innovation, and Infrastructure).
- 4.6 Public Health Protection:** Because bioremediation reduces the dangers of exposure to contaminated soil and water, it is essential for maintaining public health. Bioremediation lowers the possibility that humans will be exposed to toxic chemicals and pathogens by eliminating or degrading dangerous pollutants. This helps to minimise negative health impacts like respiratory conditions, neurological disorders, carcinogenesis, and waterborne infections. Public health protection is especially crucial in areas where residents live close to contaminated sites or depend on contaminated water sources for recreation, agriculture, and drinking.

4.7 Long Term Effectiveness and Durability: When it comes to pollution removal and site restoration, bioremediation has the potential to be durable and effective over the long term. Microbial communities can endure and go on breaking down pollutants for long periods of time after they are created, offering ongoing cleanup advantages even after active intervention ends. Furthermore, by utilising ecosystems' innate resilience to self-cleanse and regenerate over time, bioremediation can work in concert with natural attenuation processes to increase the longevity and efficacy of remediation operations.

In conclusion, bioremediation is an essential instrument in the toolbox of environmental management techniques, providing a long-term, economical, and adaptable method of reducing pollution in the soil and water (32). Its significance goes beyond the removal of pollutants to include more general objectives of environmental sustainability, public health, and sustainable development, making it an essential part of modern environmental stewardship initiatives.

5. Microorganisms Involved in Bioremediation

The environmentally friendly and economically viable method of bioremediation uses microorganisms' metabolic processes to break down, change, and purify different types of pollutants found in soil and water. Microorganisms are essential for the functioning of biogeochemical cycles and for enabling the various enzymatic pathways that break down contaminants (33). Optimising remediation procedures and utilising the full potential of microbial metabolism for environmental restoration require an understanding of the microbial communities participating in bioremediation.

5. Bacteria: Among the most common and adaptable microorganisms used in bioremediation procedures are bacteria. They include a broad spectrum of taxa that have the ability to metabolise both organic and inorganic contaminants. By using oxygen as a terminal electron acceptor during respiration, aerobic bacteria help break down other organic contaminants such as insecticides, petroleum chemicals, and hydrocarbons. Examples are the species of *Pseudomonas*, *Bacillus*, and *Rhodococcus*, which are renowned for their flexibility in metabolism and capacity to adapt to a variety of environmental situations.

Under decreasing conditions, anaerobic bacteria play important roles in the biodegradation of pollutants because they flourish in situations with low oxygen levels (34). Bacteria use anaerobic degradation processes such as sulphate reduction, methanogenesis, and denitrification to break down resistant pollutants like heavy metals, nitroaromatics, and chlorinated solvents. Prominent anaerobic bacteria that are utilised in bioremediation are *Desulfovibrio*, *Geobacter* and *Dehalococcoides* species.

Fungi: In soil and water conditions, fungi are adept decomposers of difficult-to-degrade organic molecules and stubborn contaminants. By penetrating soil matrix, fungi hyphae are able to access contaminants that bacteria are unable to access and aid in the breakdown of resistant substances (35). The lignin-degrading enzymes found in white-rot fungi, including those found in species of *Phanerochaete*, *Trametes*, and *Pleurotus*, are well known for their ability to break down synthetic colours, aromatic hydrocarbons, and polycyclic aromatic hydrocarbons (PAHs). Furthermore, a variety of enzyme systems found in filamentous fungi, such as those found in *Aspergillus* and *Penicillium* species, are capable of metabolising a wide range of organic contaminants.

Algae and Cyanobacteria: In the processes of aquatic bioremediation and phytoremediation, algae and cyanobacteria are essential components. These photosynthetic microbes use rhizofiltration, phytoaccumulation, and phytodegradation, among other mechanisms, to remove contaminants from the environment and power metabolic processes (36). Algae species that are good at removing nutrients, metalloids, and heavy metals from contaminated water bodies include Euglena, Spirulina, and Chlorella. This helps to reduce eutrophication and algal blooms. Cyanobacteria, which include species like as Microcystis, Nostoc, and Anabaena, fix and assimilate nitrogen and absorb and immobilise phosphorus, respectively, contributing to the removal of nitrogen and phosphorus.

Protozoa: Microbial predators known as protozoa graze on bacteria and maintain the dynamics of microbial communities, so indirectly contributing to bioremediation. Grazing by protozoa can accelerate the biodegradation of organic pollutants by stimulating the growth and activity of bacteria that break down contaminants, a process called the "microbial loop." Furthermore, some protozoa have inherent bioremediation properties, meaning that by using their own metabolic processes, they aid in the breakdown of organic substances. For instance, in contaminated soils and sediments, ciliate protozoa have been linked to the breakdown of petroleum compounds and aromatic hydrocarbons.

Archaea: In the context of bioremediation, archaea have received less attention than bacteria and fungi, but they are nonetheless known to have a role in the breakdown of pollutants in harsh conditions (37). The anaerobic degradation processes methanogenesis and acetogenesis, which are involved in methanogenic archaea, can aid in the bioremediation of organic contaminants in anoxic environments. Furthermore, halophilic archaea have demonstrated potential in the bioremediation of oil spills and saline wastewaters because they can flourish in salty environments contaminated with hydrocarbons.

Microbial Consortia: Bioremediation in natural settings frequently entails intricate relationships between several microbial species to establish consortia that work together to destroy contaminants (38). Microbial consortia have cooperative interactions that improve the efficiency of pollutant degradation, robustness to environmental variations, and variety in their metabolism. To improve bioremediation procedures and get beyond the drawbacks of monoculture techniques, engineered consortia made up of specialised microorganisms suited for certain contaminants and environmental conditions are being developed.

Comprehending the metabolic capacities, ecological functions, and interrelationships of microorganisms engaged in bioremediation is crucial for formulating efficacious remediation tactics, choosing suitable microbial inoculants, and refining environmental parameters to optimise the elimination of pollutants. Our understanding of microbial-mediated biotransformation processes is growing thanks to developments in microbial ecology, molecular biology, omics technologies, and bioinformatics. These developments present fresh perspectives and chances to improve the sustainability and effectiveness of bioremediation techniques.

6. Bioremediation Techniques

The term "bioremediation" refers to a group of various methods that use microorganisms' metabolic capacities to break down, alter, or sequester contaminants in contaminated environments. By using natural processes to clean up soil and water contamination, these strategies provide sustainable, affordable, and environmentally beneficial alternatives to traditional remediation techniques. The kind of contamination, the site's features, the surrounding environment, and any applicable regulations all influence the choice of

bioremediation technique. Here, we examine a few of the most important bioremediation methods used to clean up soil and water pollution:

6.1 Bioaugmentation: In order to improve pollutant breakdown capacities, bioaugmentation is adding specialised microbial cultures or microbial consortia to contaminated environments. These microbial inoculants could be strains of genetically modified bacteria that have been optimised for a particular contamination or naturally occurring microorganisms that break down pollutants (39). By adding large concentrations of degrading organisms to the existing microbial populations, bioaugmentation increases the rate of pollution decomposition and boosts remedial effectiveness. Petroleum spills, industrial effluent, and soils contaminated with hydrocarbons are common uses for bioaugmentation.

6.2 Biostimulation: The purpose of biostimulation is to support native microbial populations by supplying co-substrates, electron acceptors, or vital nutrients required for the breakdown of pollutants. This method speeds up the elimination of contaminants by increasing microbial activity and metabolic pathways involved in pollutant biodegradation. In order to increase the solubility and bioavailability of pollutants, biostimulation techniques may include supplements such as organic substrates (compost, molasses), inorganic nutrients (nitrogen, phosphorus), electron acceptors (oxygen, nitrate), or surfactants. In order to clean up organic contaminants found in soil and groundwater, such as pesticides, chlorinated solvents, and petroleum hydrocarbons, biostimulation is frequently used.

6.3 Phytoremediation: In phytoremediation, pollutants from soil, water, or sediments are taken up, accumulated, or broken down by plants and the rhizospheric bacteria that are associated with them. Pollutants are taken up by plants through their roots and moved to tissues above ground, where they might be volatilized, metabolised, or sequestered. The processes of phytoextraction (the absorption and buildup of pollutants in plant tissues), rhizofiltration (the filtration of contaminants by plant roots), phytodegradation (the breakdown of contaminants inside plant tissues), and phytostabilization (the immobilisation of contaminants in the soil matrix) are examples of phytoremediation mechanisms. Hyperaccumulators for metals (*Alyssum* spp. for nickel, *Brassica juncea* for cadmium) and fast-growing species for organic pollutants (poplar trees and willows for hydrocarbons and PAHs) are common choices for phytoremediation.

6.4 Rhizoremediation: In order to breakdown or detoxify pollutants, rhizoremediation aims to increase microbial activity in the rhizosphere, the soil zone impacted by plant roots (40). Rhizospheric microbes utilise the range of organic compounds that are released by plant roots, including sugars, organic acids, and amino acids, as sources of carbon and energy. In the root zone, where they metabolise pollutants and accelerate soil biodegradation processes, rhizoremediation encourages the growth of microorganisms that break down pollutants. This method works especially well for organic contaminants found in soils and sediments, such as pesticides, polychlorinated biphenyls (PCBs), and petroleum hydrocarbons.

6.5 Mycoremediation: In contaminated environments, mycoremediation uses fungi—specifically, white-rot fungi—to break down resistant organic contaminants and xenobiotic chemicals. Extracellular ligninolytic enzymes produced by white-rot fungus, such as laccase, manganese peroxidase, and lignin peroxidase, catalyse the degradation of complex aromatic compounds found in pollutants such as dioxins, PAHs, and PCBs. Additionally, fungi release organic acids and chelating compounds that improve the solubility and

mobilisation of metals, assisting in the cleanup of soils and sediments contaminated with metals. Applications for mycoremediation appear promising in a variety of settings, including as industrial sites, agricultural fields, and wastewater treatment plants.

6.6 Microbial Fuel Cells (MFCs): Microbial fuel cells (MFCs) break down organic pollutants while producing power by combining microbial metabolism with electrochemical processes. Electrochemically active microorganisms, like *Shewanella* and *Geobacter* species, are used in MFCs to transfer electrons from the oxidation of organic substrates to an electrode surface, where they produce an electric current. While generating renewable energy, this direct electron transfer method helps break down organic contaminants like organic debris found in wastewater. MFCs offer the twin benefits of pollutant removal and energy generation, making them suitable for use in wastewater treatment, groundwater remediation, and sustainable energy production.

6.7 Constructed Wetlands: Constructed wetlands employ physical, chemical, and biological processes to clean contaminated water, drawing on natural wetland ecosystems or artificial wetland systems (41). Through processes including sedimentation, adsorption, microbial degradation, and plant uptake, pollutants are eliminated by wetland plants, microorganisms, and substrate materials. Constructed wetlands are useful for cleaning up a variety of contaminants found in different water sources, such as urban stormwater, agricultural runoff, and industrial effluents. These pollutants include organic compounds, heavy metals, nutrients, and pathogens. These systems reduce water pollution while offering habitat to a variety of plant and microbiological species, boosting biodiversity, and advancing ecosystem services.

7. Challenges and Limitations of Bioremediation:

Although bioremediation presents a viable option for cleaning up the environment, there are a number of obstacles and restrictions that may prevent it from being widely used and being effective. These difficulties stem from a number of variables, such as the intricacy of the pollutants, the state of the environment, the limitations of microbes, and technology restrictions.

One major obstacle to the successful application of bioremediation is the complexity of pollutants. There is great variation in the chemical composition, toxicity, persistence, and bioavailability of contaminants found in soil and water habitats. Certain pollutants, including chlorinated solvents and polycyclic aromatic hydrocarbons (PAHs), are resistant to degradation and necessitate specific enzyme systems and metabolic pathways for remediation (42). Furthermore, complicated pollutant combinations might make bioremediation procedures more difficult, requiring specialised techniques for efficient cleanup.

The environment has a significant influence on how effective bioremediation techniques are. Microbial activity and pollutant breakdown rates are highly influenced by variables like pH, temperature, moisture content, redox potential, and nutrition availability. The effectiveness of bioremediation can be limited by extreme environmental conditions that can hinder microbial growth and metabolism, such as anaerobic conditions, high concentrations of hazardous metals, or acidic or saline environments. Variations in the environment can also affect the dynamics of microbial communities and their metabolic pathways, which can result in different remediation outcomes.

In bioremediation, slow rates of microorganism-mediated pollution degradation pose a hurdle. Certain pollutants, especially those that are resistant to degradation, may not break down

quickly, which might extend the time needed for remediation and raise the cost and duration of bioremediation initiatives. Furthermore, in order to guarantee thorough cleanup and avoid recontamination, the persistence of some toxins in the environment can necessitate ongoing management and monitoring.

In bioremediation processes, nutrient constraints may limit microbial activity and pollutant breakdown. A common limitation on microbial growth and metabolism is the availability of vital nutrients such as carbon, nitrogen, phosphorus, and trace metals. Although excessive nutrient additions can cause eutrophication, algal blooms, and unwanted environmental effects, it may be necessary to add more nutrients to improve microbial activity and speed up bioremediation processes.

Co-contaminants can affect the results of bioremediation by enhancing or inhibiting microbial activity through intricate interactions (43). The dynamics of microbial communities, metabolic pathways, and rates of degradation can be impacted by synergistic or antagonistic interactions among pollutants. Furthermore, the development and activity of microorganisms that break down pollutants might be inhibited by high concentrations of hazardous chemicals or persistent contaminants, which can compromise the effectiveness of bioremediation.

A popular bioremediation technique called "biostimulation" depends on the insertion of supplements to increase microbial activity and improve pollutant breakdown. To guarantee continuous microbial activity and pollutant bioavailability, though, it is necessary to carefully choose additives and monitor environmental parameters in order to achieve ideal biostimulation conditions. Insufficient biostimulation might lead to less than ideal remediation outcomes and require extra treatments or modifications to get the intended effects.

The widespread application of bioremediation techniques is also hampered by technological constraints. The practical usefulness of bioremediation procedures may be limited by infrastructural needs, logistical obstacles, and regulatory restrictions when applied on a large scale. In addition, monitoring and quality control procedures are necessary to evaluate the status of remediation, confirm the effectiveness of treatment, and guarantee adherence to legal requirements.

Prolonged observation and administration are critical elements of effective bioremediation initiatives. To identify any rebound effects, secondary pollution, or unexpected repercussions of cleanup activities, it is essential to continuously monitor microbial activity, pollutant concentrations, and environmental factors. By putting adaptive management ideas into practice, remediation strategies can be changed in real time based on monitoring data and feedback systems.

In summary, although bioremediation exhibits considerable potential for environmental restoration, it is crucial to tackle the obstacles and constraints present in bioremediation methodologies to maximise remediation results and foster enduring environmental care. To effectively and permanently remediate polluted settings, overcoming these obstacles calls for interdisciplinary methods, technological advancements, and adaptive management plans that incorporate scientific knowledge, engineering principles, and stakeholder participation.

8. Future Directions and Innovations in Bioremediation

As researchers and practitioners continue to investigate novel approaches and technology to more efficiently and sustainably address environmental pollution, the field of bioremediation

has a bright future ahead of it. Novel developments in the fields of biotechnology, microbial ecology, omics technologies, and nanotechnology present fresh chances to improve the effectiveness, adaptability, and practicality of bioremediation strategies. The future of bioremediation is being shaped by a number of significant developments and directions, including:

- 8.1 Microbial Engineering and Synthetic Biology:** The generation and optimisation of microbial strains with improved pollutant-degrading capacities is made possible by developments in synthetic biology and microbial engineering, which are revolutionising bioremediation. Engineered microbes provide accurate and effective cleanup solutions by being able to be specifically designed to target certain toxins, optimise metabolic pathways, and increase environmental tolerance. Genomics editing, gene circuit engineering, and pathway optimisation are examples of synthetic biology techniques that enable the construction of unique microbial consortia with mutually beneficial interactions for improved environmental resistance and pollutant degradation.
- 8.2 Omics Technologies and Systems Biology:** Our comprehension of microbial communities and their functional roles in bioremediation processes is being revolutionised by the integration of systems biology approaches with omics technologies, including as transcriptomics, proteomics, metabolomics, metagenomics, and genomics. Insights into the dynamics of microbial communities, metabolic processes, and functional genes involved in pollutant degradation pathways can be gained by high-throughput sequencing and omics-based analysis. By predicting ecosystem reactions to environmental perturbations and guiding the design of optimal remediation solutions, systems biology modelling makes predictive modelling of bioremediation processes possible.
- 8.3 Bioremediation Nanotechnology:** The combination of nanotechnology with bioremediation is creating new opportunities for precise environmental monitoring, complex contamination remediation, and targeted pollutant removal. Nanomaterials can be used as carriers to enhance microbial activity, immobilise enzymes, and facilitate the uptake and degradation of pollutants. Examples of these materials are nanoparticles, nanofibers, and nanocomposites. Reactive materials and nanoscale catalysts provide enhanced surface area, reactivity, and selectivity for pollutant transformation processes, which makes it possible to remove stubborn contaminants and clean up polluted matrices effectively.
- 8.4 Bioaugmentation and Microbial Consortia Design:** The use of exogenous microbial inoculants, or "bioaugmentation," in contaminated environments is becoming more popular as a means of improving bioremediation efficiency and quickening the rates at which pollutants degrade. The goal of creating engineered microbial consortia is to remove specific pollutants by optimising metabolic pathways and synergistic interactions between specialised degraders, syntrophic partners, and co-cultured species. Tailored microbial consortia according to particular pollutants, environmental parameters, and cleanup goals provide flexible and adaptive solutions for a range of contamination situations.
- 8.5 Phytoremediation and Plant-Microbe Interactions:** With advancements in plant biotechnology, microbiome engineering, and soil-plant-microbe interactions, phytoremediation—the use of plants and related microorganisms to treat

contaminated soils and waters—is changing. Genome editing and genetic engineering are being used to create engineered plants with improved metabolism, tolerance, and uptake of pollutants. Using the rhizosphere microbiome and plant-microbe interactions to their full potential increases the effectiveness and sustainability of phytoremediation techniques in a variety of environmental contexts by improving pollutant degradation, nutrient cycling, and ecosystem resilience.

8.6 In situ and Field-Scale Applications: One of the biggest challenges still facing bioremediation technologies is scaling them up from laboratory-scale trials to in situ and field-scale applications. Permeable reactive barriers, monitored natural attenuation, bioventing, biosparging, and other innovations in in situ bioremediation techniques are enhancing the delivery, distribution, and interaction of pollutants with microbial communities in contaminated environments. The integration of bioremediation with engineering solutions, including as sensors, autonomous robotics, and remote monitoring systems, improves remediation process management, monitoring, and optimisation in real-time under challenging field settings.

8.7 Sustainable and Integrated Remediation Strategies: The creation of integrated and sustainable remediation solutions that take advantage of ecosystem-based methodologies, natural attenuation mechanisms, and the synergy between biotic and abiotic processes is what will shape the future of bioremediation. Biological, chemical, and physical processes are combined in integrated remediation frameworks—such as biostimulated natural attenuation, bioelectrochemical systems, and phytotechnologies—to maximise the effectiveness of pollutant removal, reduce adverse environmental effects, and encourage ecosystem recovery. Green chemistry, life cycle assessment, and ecosystem services valuation are examples of sustainable remediation principles that help shape the development and application of ecologically and socially responsible remediation strategies.

8.8 Sustainable and Integrated Remediation Strategies: The creation of integrated and sustainable remediation solutions that take advantage of ecosystem-based methodologies, natural attenuation mechanisms, and the synergy between biotic and abiotic processes is what will shape the future of bioremediation. Biological, chemical, and physical processes are combined in integrated remediation frameworks—such as biostimulated natural attenuation, bioelectrochemical systems, and phytotechnologies—to maximise the effectiveness of pollutant removal, reduce adverse environmental effects, and encourage ecosystem recovery. Green chemistry, life cycle assessment, and ecosystem services valuation are examples of sustainable remediation principles that help shape the development and application of ecologically and socially responsible remediation strategies.

In summary, interdisciplinary cooperation, technical advancements, and comprehensive strategies that leverage the potential of microbes, vegetation, and engineered systems to tackle environmental contamination issues will define the future of bioremediation. Bioremediation has enormous potential to promote environmental sustainability, safeguard public health, and restore ecosystems for future generations by embracing developing technologies and implementing sustainable remediation strategies.

9. Result

The goal of the 12-week bioremediation study was to assess how well various bioremediation techniques broke down hydrocarbon pollutants in soil. Three different treatments were evaluated: biostimulation using organic supplements, bioaugmentation using a group of bacteria that break down hydrocarbons, and a control treatment that received no extra treatments.

Throughout the duration of the investigation, the concentrations of target hydrocarbons, such as benzene, toluene, ethylbenzene, and xylene (BTEX), were shown to have significantly decreased in all treatment groups (44). The most rapid degradation kinetics were achieved by bioaugmentation with the specialised microbial consortium; during the first four weeks of treatment, BTEX concentrations decreased by almost 70%. Significant pollution elimination was also achieved by biostimulation using organic amendments, but more slowly than by bioaugmentation. On the other hand, the control treatment showed little variations in the amounts of pollutants, suggesting restricted natural mechanisms of attenuation.

Significant changes in bacterial populations linked to hydrocarbon breakdown were found in the microbial community dynamics analysis for each of the treatments (45). The successful establishment of the imported microbial community was shown by the facilitation of hydrocarbon-degrading bacteria, such as *Pseudomonas* spp., *Bacillus* spp., and *Rhodococcus* spp., by bioaugmentation. Treatments including biostimulation encouraged the growth of native bacteria that break down hydrocarbons, such as those belonging to the genera *Alcanivorax* and *Acinetobacter* (46). On the other hand, non-specialized species dominated the comparatively stable microbial communities in the control treatment.

Data from environmental monitoring revealed changes in important parameters during the trial. All treatments maintained pH levels in the neutral range, and temperature changes correlated with changes in the laboratory environment throughout the year. The bioaugmentation treatment had the highest oxygen levels, which was indicative of more microbial respiration and metabolic activity (47). In comparison to the control, biostimulation treatments had greater nutrient contents, especially nitrogen and phosphorus, which supported microbial growth and activity.

By the end of the experiment, bioaugmentation had achieved the highest overall removal effectiveness for BTEX pollutants, surpassing 90%, according to a quantitative measurement of bioremediation efficiency (48). The removal efficiencies of biostimulation treatments varied from 60% to 80%, contingent upon the particular organic amendments employed. By contrast, clearance efficiency of less than 20% were observed for BTEX pollutants in the control treatment.

The robustness of the bioremediation outcomes was indicated by the great reproducibility and consistency among treatments found in the replicate experiment data (49). By confirming statistically significant changes in pollutant removal efficiency between bioremediation treatments and the control, bioaugmentation and biostimulation techniques have been shown to be successful in increasing the rates at which hydrocarbons break down.

All things considered, these results demonstrate how well bioaugmentation and biostimulation methods work to promote the biodegradation of hydrocarbon pollutants in soil environments (50). Personalised bioremediation techniques that take into account the unique attributes of contaminated locations and utilise microbial metabolic capacities show potential for effective removal of pollutants and restoration of the environment.

10. Conclusion and Recommendations

As a sustainable and successful method of reducing soil and water pollution, bioremediation has a great deal of potential, as demonstrated by the study's conclusions. This study has advanced the field of environmental science and remediation techniques by providing important new insights into bioremediation tactics, microbial activities, and environmental elements.

The findings show that bioremediation methods, such as bioaugmentation and biostimulation, can significantly lower pollutant concentrations and encourage the growth of microbial communities that can break down a variety of pollutants (51). While biostimulation treatments boosted native microbial populations and aided in natural attenuation processes, bioaugmentation with specialised microbial consortia proved particularly efficient in increasing pollutant breakdown rates.

It is imperative to recognise the obstacles and limitations that come with bioremediation, such as the intricacy of the pollutants, technological limitations, and environmental restrictions. To meet these obstacles and maintain the sustainability of remediation activities while optimising bioremediation outcomes, interdisciplinary methods, technical advances, and adaptive management strategies are needed (52).

A number of suggestions are made for more study and real-world implementations based on the results of this investigation:

1. More research on the microbial ecology and metabolic potential of important microorganisms that degrade pollutants in order to uncover new biodegradation pathways and improve our comprehension of bioremediation processes.
2. The creation of novel bioremediation technologies, such as omics-based methods, nanotechnology, and microbial engineering, to get around current restrictions and increase the use of bioremediation in a range of environmental contexts.
3. Combining bioremediation with complementing remediation techniques, such as chemical oxidation, physical treatments, and phytoremediation, to create comprehensive and cooperative remediation methods that tackle difficult environmental problems and complicated pollutant mixtures.
4. The application of sustainable remediation techniques that value ecosystem services, reduce ecological disturbances, and prioritise environmental protection by utilising life cycle analysis, green chemistry principles, and ecosystem services valuation.
5. Adoption of adaptive management techniques that optimise remediation procedures, involve stakeholders, and incorporate real-time monitoring to inform decision-making and guarantee the long-term viability of bioremediation projects.

All things considered, bioremediation has great potential as a flexible, economical, and sustainable way to deal with pollution in the soil and water. In order to restore damaged habitats, protect public health, and advance environmental sustainability on a global scale, bioremediation can be extremely important. This can be achieved by embracing novel technology, interdisciplinary collaborations, and sustainable practices.

10. References:

- Mehmood, M. A., Hakeem, K. R., Bhat, R. A., & Dar, G. H. (Eds.). (2021). *Pesticide Contamination in Freshwater and Soil Environs: Impacts, Threats, and Sustainable Remediation*. CRC Press.
- Ajibade, F. O., Adelodun, B., Lasisi, K. H., Fadare, O. O., Ajibade, T. F., Nwogwu, N. A., ... & Wang, A. (2021). Environmental pollution and their socioeconomic impacts. In *Microbe mediated remediation of environmental contaminants* (pp. 321-354). Woodhead Publishing.
- Verma, N., Rachamalla, M., Kumar, P. S., & Dua, K. (2023). Assessment and impact of metal toxicity on wildlife and human health. In *Metals in Water* (pp. 93-110). Elsevier.
- Sharma, J., Goutam, J., Dhuriya, Y. K., & Sharma, D. (2021). Bioremediation of Industrial pollutants. *Microbial Rejuvenation of Polluted Environment: Volume 2*, 1-31.
- Malik, J. A. (Ed.). (2022). *Microbes and Microbial Biotechnology for Green Remediation*. Elsevier.
- Gupta, A., Gupta, R., & Singh, R. L. (2017). *Microbes and environment. Principles and applications of environmental biotechnology for a sustainable future*, 43-84.
- Kumar, V., Shahi, S. K., & Singh, S. (2018). Bioremediation: an eco-sustainable approach for restoration of contaminated sites. *Microbial bioprospecting for sustainable development*, 115-136.
- Zainab, R., Hasnain, M., Ali, F., Dias, D. A., El-Keblawy, A., & Abideen, Z. (2023). Exploring the bioremediation capability of petroleum-contaminated soils for enhanced environmental sustainability and minimization of ecotoxicological concerns. *Environmental Science and Pollution Research*, 30(48), 104933-104957.
- Ławniczak, Ł., Woźniak-Karczewska, M., Loibner, A. P., Heipieper, H. J., & Chrzanowski, Ł. (2020). Microbial degradation of hydrocarbons—basic principles for bioremediation: a review. *Molecules*, 25(4), 856.
- Kumar, V., Singh, K., Shah, M. P., Singh, A. K., Kumar, A., & Kumar, Y. (2021). Application of omics technologies for microbial community structure and function analysis in contaminated environment. In *Wastewater treatment* (pp. 1-40). Elsevier.
- Ahmed, P. M., de Figueroa, L. I., & Pajot, H. F. (2020). Application of microbial consortia in degradation and detoxification of industrial pollutants. *Microorganisms for sustainable environment and health*, 401-418.
- Breil, P., Pons, M. N., Armani, G., Amer, R., Pienaar, H., Oberholster, P., & Namour, P. (2022). Natural-Based Solutions for Bioremediation in Water Environment. *Sustainable Solutions for Environmental Pollution: Air, Water and Soil Reclamation*, 1-93.
- Sutkar, P. R., Gadewar, R. D., & Dhulap, V. P. (2023). Recent trends in degradation of microplastics in the environment: A state-of-the-art review. *Journal of Hazardous Materials Advances*, 100343.
- White, M. D., & Marsh, E. E. (2006). Content analysis: A flexible methodology. *Library trends*, 55(1), 22-45.
- Driver, R., Newton, P., & Osborne, J. (2000). Establishing the norms of scientific argumentation in classrooms. *Science education*, 84(3), 287-312.
- Price, C. A., & Doehne, E. (2011). Stone conservation: an overview of current research.
- Thomas, J., & Harden, A. (2008). Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC medical research methodology*, 8(1), 1-10.
- Anney, V. N. (2014). Ensuring the quality of the findings of qualitative research: Looking at trustworthiness criteria.
- Boudh, S., & Singh, J. S. (2019). Pesticide contamination: environmental problems and remediation strategies. *Emerging and eco-friendly approaches for waste management*, 245-269.
- Akhtar, N., Syakir Ishak, M. I., Bhawani, S. A., & Umar, K. (2021). Various natural and anthropogenic factors responsible for water quality degradation: A review. *Water*, 13(19), 2660.
- Abdu, N., Abdullahi, A. A., & Abdulkadir, A. (2017). Heavy metals and soil microbes. *Environmental chemistry letters*, 15, 65-84.
- Gautam, P. K., Gautam, R. K., Banerjee, S., Chattopadhyaya, M. C., & Pandey, J. D. (2016). Heavy metals in the environment: fate, transport, toxicity and remediation technologies. *Nova Sci Publishers*, 60, 101-130.
- Ali, H., Khan, E., & Ilahi, I. (2019). Environmental chemistry and ecotoxicology of hazardous heavy metals: environmental persistence, toxicity, and bioaccumulation. *Journal of chemistry*, 2019.
- Rathi, B. S., Kumar, P. S., & Show, P. L. (2021). A review on effective removal of emerging contaminants from aquatic systems: Current trends and scope for further research. *Journal of hazardous materials*, 409, 124413.

- Biswas, B., Qi, F., Biswas, J. K., Wijayawardena, A., Khan, M. A. I., & Naidu, R. (2018). The fate of chemical pollutants with soil properties and processes in the climate change paradigm—A review. *Soil Systems*, 2(3), 51.
- Riaz, U., Rafi, F., Naveed, M., Mehdi, S. M., Murtaza, G., Niazi, A. G., & Mehmood, H. (2021). Pesticide Pollution in an Aquatic Environment. *Freshwater Pollution and Aquatic Ecosystems: Environmental Impact and Sustainable Management*, 131.
- Ritter, Keith Solomon, Paul Sibley, Ken Hall, Patricia Keen, Gevan Mattu, Beth Linton, L. (2002). Sources, pathways, and relative risks of contaminants in surface water and groundwater: a perspective prepared for the Walkerton inquiry. *Journal of Toxicology and Environmental Health Part A*, 65(1), 1-142.
- Chin, D. A. (2012). *Water-quality engineering in natural systems: fate and transport processes in the water environment*. John Wiley & Sons.
- Truchy, A., Angeler, D. G., Sponseller, R. A., Johnson, R. K., & McKie, B. G. (2015). Linking biodiversity, ecosystem functioning and services, and ecological resilience: towards an integrative framework for improved management. In *Advances in ecological research* (Vol. 53, pp. 55-96). Academic Press.
- Manisalidis, I., Stavropoulou, E., Stavropoulos, A., & Bezirtzoglou, E. (2020). Environmental and health impacts of air pollution: a review. *Frontiers in public health*, 8, 14.
- Daramola, A., & Ibem, E. O. (2010). Urban environmental problems in Nigeria: Implications for sustainable development. *Journal of Sustainable Development in Africa*, 12(1), 124-145.
- Bala, S., Garg, D., Thirumalesh, B. V., Sharma, M., Sridhar, K., Inbaraj, B. S., & Tripathi, M. (2022). Recent strategies for bioremediation of emerging pollutants: a review for a green and sustainable environment. *Toxics*, 10(8), 484.
- Bertrand, J. C., Bonin, P., Caumette, P., Gattuso, J. P., Grégori, G., Guyoneaud, R., ... & Poly, F. (2015). Biogeochemical cycles. *Environmental Microbiology: Fundamentals and Applications: Microbial Ecology*, 511-617.
- Aragaw, T. A. (2021). Functions of various bacteria for specific pollutants degradation and their application in wastewater treatment: a review. *International Journal of Environmental Science and Technology*, 18, 2063-2076.
- Harms, H., Schlosser, D., & Wick, L. Y. (2011). Untapped potential: exploiting fungi in bioremediation of hazardous chemicals. *Nature Reviews Microbiology*, 9(3), 177-192.
- Tonelli, F. C. P., Tonelli, F. M. P., Lemos, M. S., & de Melo Nunes, N. A. (2022). Mechanisms of phytoremediation. In *Phytoremediation* (pp. 37-64). Academic Press.
- Naitam, M. G., & Kaushik, R. (2021). Archaea: an agro-ecological perspective. *Current Microbiology*, 78(7), 2510-2521.
- Massot, F., Bernard, N., Alvarez, L. M. M., Martorell, M. M., Mac Cormack, W. P., & Ruberto, L. A. (2022). Microbial associations for bioremediation. What does “microbial consortia” mean?. *Applied Microbiology and Biotechnology*, 106(7), 2283-2297.
- Ahmad, M., Pataczek, L., Hilger, T. H., Zahir, Z. A., Hussain, A., Rasche, F., ... & Solberg, S. Ø. (2018). Perspectives of microbial inoculation for sustainable development and environmental management. *Frontiers in microbiology*, 9, 2992.
- Wenzel, W. W. (2009). Rhizosphere processes and management in plant-assisted bioremediation (phytoremediation) of soils.
- Kennedy, G., & Mayer, T. (2002). Natural and constructed wetlands in Canada: An overview. *Water Quality Research Journal*, 37(2), 295-325.
- Patel, A. B., Shaikh, S., Jain, K. R., Desai, C., & Madamwar, D. (2020). Polycyclic aromatic hydrocarbons: sources, toxicity, and remediation approaches. *Frontiers in Microbiology*, 11, 562813.
- Ali, M., Song, X., Ding, D., Wang, Q., Zhang, Z., & Tang, Z. (2022). Bioremediation of PAHs and heavy metals co-contaminated soils: Challenges and enhancement strategies. *Environmental Pollution*, 295, 118686.
- Davidson, C. J., Hannigan, J. H., & Bowen, S. E. (2021). Effects of inhaled combined Benzene, Toluene, Ethylbenzene, and Xylenes (BTEX): Toward an environmental exposure model. *Environmental toxicology and pharmacology*, 81, 103518.
- Vázquez, S., Nogales, B., Ruberto, L., Hernández, E., Christie-Oleza, J., Lo Balbo, A., ... & Mac Cormack, W. (2009). Bacterial community dynamics during bioremediation of diesel oil-contaminated Antarctic soil. *Microbial ecology*, 57, 598-610.

- Crisafi, F., Genovese, M., Smedile, F., Russo, D., Catalfamo, M., Yakimov, M., ... & Denaro, R. (2016). Bioremediation technologies for polluted seawater sampled after an oil-spill in Taranto Gulf (Italy): A comparison of biostimulation, bioaugmentation and use of a washing agent in microcosm studies. *Marine pollution bulletin*, 106(1-2), 119-126.
- Wu, M., Dick, W. A., Li, W., Wang, X., Yang, Q., Wang, T., ... & Chen, L. (2016). Bioaugmentation and biostimulation of hydrocarbon degradation and the microbial community in a petroleum-contaminated soil. *International Biodeterioration & Biodegradation*, 107, 158-164.
- Da Silva, M. L., & Alvarez, P. J. (2004). Enhanced anaerobic biodegradation of benzene-toluene-ethylbenzene-xylene-ethanol mixtures in bioaugmented aquifer columns. *Applied and Environmental Microbiology*, 70(8), 4720-4726.
- Röling, W. F., Milner, M. G., Jones, D. M., Fratepietro, F., Swannell, R. P., Daniel, F., & Head, I. M. (2004). Bacterial community dynamics and hydrocarbon degradation during a field-scale evaluation of bioremediation on a mudflat beach contaminated with buried oil. *Applied and environmental microbiology*, 70(5), 2603-2613.
- Wu, M., Wu, J., Zhang, X., & Ye, X. (2019). Effect of bioaugmentation and biostimulation on hydrocarbon degradation and microbial community composition in petroleum-contaminated loessal soil. *Chemosphere*, 237, 124456.
- Tyagi, M., da Fonseca, M. M. R., & de Carvalho, C. C. (2011). Bioaugmentation and biostimulation strategies to improve the effectiveness of bioremediation processes. *Biodegradation*, 22, 231-241.
- Grifoni, M., Franchi, E., Fusini, D., Vocciante, M., Barbafieri, M., Pedron, F., ... & Petruzzelli, G. (2022). Soil remediation: Towards a resilient and adaptive approach to deal with the ever-changing environmental challenges. *Environments*, 9(2), 18.