

Work Engagement And Occupational Stress In Nurses And Other Healthcare Workers: The Role Of Organizational And Personal Factors

Norah Hamoud Abeer Alanazi¹, Eidah Hamoud A Alanazi², Amal lahawh alonazi³, Albandari Faleh Aldusaeri⁴, Nouf Saud Ali Alrowdan⁵, Latifa Ayad Fahd Al-Harbi⁶, Rasha Salem Alanazi⁷, Abdullah Mobark Ali Alsahli⁸

Abstract

The aim of this paper was to review the literature on factors related to role stress in nurses, and present strategies for addressing this issue based on the findings of this review while considering potential areas for development and research. Computerized databases were searched as well as hand searching of articles in order to conduct this review. This review identified multiple factors related to the experience of role stress in nurses. Role stress, in particular, work overload, has been reported as one of the main reasons for nurses leaving the workforce. This paper concludes that it is a priority to find new and innovative ways of supporting nurses in their experience of role stress. Some examples discussed in this article include use of stress education and management strategies; team-building strategies; balancing priorities; enhancing social and peer support; flexibility in work hours; protocols to deal with violence; and retention and attraction of nursing staff strategies. These strategies need to be empirically evaluated for their efficacy in reducing role stress.

Key words *age and role stress, role stress and nurses, role stress and staffing, role stress and violence, role stress reduction.*

INTRODUCTION

Role stress in nurses continues to be an area of great interest to the profession, particularly as stress affects the mental and physical health of the nurses as well as having an economic cost to the community (Cooper, 1998). Evidence by a review of the literature conducted by Lambert and Lambert (2001) found that over 100 articles on role stress in nursing had been published in English over the past two decades, signifying the underlying importance of this area and its continued lack of resolution. Changes continue to occur rapidly in the health care area, and currently an epidemic exists of an international shortage of nursing staff that shows little sign of abating (Janiszewski Goodin, 2003). In the USA, more than 10 000 nurses are required to meet existing shortages, with demand outstripping supply by more than 800 000 by 2020 (Buerhaus, 2004). In Australia, the shortage is the greatest since World War II and there will be a shortage of up to 40 000 nurses by 2010 (RMIT University, 2003).

¹Nursing technician, Southern Nazim Health Center, Riyadh.

²Nursing technician, Qaisumah General Hospital, Hafar Al-Batin.

³Nurse technician, Qurtopa PHC, Riyadh.

⁴Nurse technician, Community health and public health, Riyadh.

⁵Nurse Technician, Community and Public Health, Riyadh.

⁶Nursing technician, Qaisumah Hospital, Hafar Al-Batin.

⁷Nursing, Granada phc, Riyadh.

⁸Nurse technician, Alyatma health centre, Almadinah city.

This situation, coupled with an environment of increasing consumption of health care goods and services, indicates the need for continued attention in identifying the most problematic areas for nurses in terms of work stress. This information can be used in supporting nurses in their roles.

This article reviews literature on factors related to role stress in nurses from 2000 to 2003, followed by a discussion on potential suggestions for strategies to reduce this experience for nurses, based on the literature.

SEARCH STRATEGY

A recent review of the literature was conducted using computerized databases that were searched over the time period 2000–2003 and included CINAHL (nursing and allied health), PsychINFO (psychology) and SSCI (the Social Science Citation Index). Hand searching of articles was also performed as well as citations in papers identified by the above searches. Key words included ‘role stress’ and ‘nurs*’, and ‘nurs* burnout’.

This search resulted in approximately 500 articles, of which 68 were relevant to this paper.

Definitions of stress, role stress and burnout

The experience of stress occurs when situations are perceived as exceeding one’s resources (Lazarus & Folkman, 1984). Work stressors are ‘antecedent conditions within one’s job or the organization which require adaptive responses on the part of the employees’ (Jex & Beehr, 1991 p. 312). Role stress may be viewed as the consequence of disparity between an individual’s perception of the characteristics of a specific role and what is actually being achieved by the individual currently performing the specific role (Lambert & Lambert, 2001). Thus, when role expectations exceed what is achieved, role stress occurs. Burnout is a role stress reaction and may be regarded as a syndrome of emotional exhaustion, depersonalization and reduced personal accomplishment (Maslach & Jackson, 1986).

REVIEW OF THE LITERATURE ON ROLE STRESS AND NURSING

A review comparing international research on role stress and strain conducted by Lambert and Lambert (2001) found that most of the studies in this field have been conducted in the USA and the UK. The key areas identified by the studies can be grouped as work environment factors, followed by the next most common aspect studied being the factors which influence and predict role stress. In smaller proportions, physiological and attitudinal factors and evaluation of instrumentation were also studied.

Common work environment factors found to be associated with role stress were: having little control in one’s job; high job demands and low supportive relationships (e.g. Chapman, 1993; Fong, 1993; Glass et al., 1993; Tovey & Adams, 1999; Webster & Hackett, 1999; Cheng et al., 2000); dealing with death and dying; being moved among different patient care units within the organization; shortage of essential resources, including nursing staff; and work overload (Foxall et al., 1990; Frisch et al., 1991; Snape & Cavanagh, 1993; Hatcher & Laschinger, 1996; Lally & Pearce, 1996; Avallone & Gibbon, 1998; Chung & Corbett, 1998; Murray, 1998). Environmental factors highlighted by the studies included uncooperative family members and clients, inability to reach physicians and unfamiliarity with situations (e.g. Walcott-McQuigg & Ervin, 1992); concern for poor quality of nursing staff, medical staff and patient care

(Scalzi, 1990); inability to deliver quality nursing care (Boswell, 1992); shift rotation (Robinson & Lewis, 1990); time demands and state laws restricting the ability to carry out the advanced practice role (Manderino et al., 1994); poor relationships with supervisors, coworkers and physicians (Decker, 1997); and low organizational commitment (Lee & Henderson, 1996).

Given that nursing shortages are at an all time high it is important that a more recent review of the literature on role stress and factors related to the experience be conducted, as well as suggestions for strategies to reduce role stress among nurses. This review also aims to obtain a wider international perspective as most studies have focused on the USA/UK experience.

RELATIONSHIP BETWEEN ROLE STRESS AND NURSING SHORTAGES

Janiszewski Goodin (2003) conducted a recent literature review of the nursing shortage in the USA. Nursing is characterized by an increasingly older group of workers as well as an undersupply of employees. The review identified four main areas contributing to the nursing shortage: the aging registered nurse (RN) workforce; declining enrollment in undergraduate nursing programs; the changing work climate; and the associated 'poor' image of nursing portrayed in the community. The work climate is characteristically that of more individuals with acute and chronic conditions requiring more complex care, contrasted by shorter hospitalization stays requiring RNs to care for and stabilize patients in a shorter time period. This situation leads to a significant increase in role stress. The American Nurses Association (ANA) (2001a) posted a national nursing survey online in 2000. Nearly 7300 nurses completed the staffing and patient care survey. Quantitative data revealed that the increased patient load and decreased time to provide direct patient care resulted in less satisfaction in the nurses' jobs. In addition, 40–60% of respondents reported that they frequently skipped meals and breaks to care for patients, felt increased pressure to accomplish their work, and participated in mandatory overtime. These factors contributed to increased dissatisfaction with their work environment and a lack of capacity to care for self. Although the large sample size lends validity to the findings, the self-report nature of this study may have biased findings.

Aiken et al. (2001) conducted a study of 700 hospitals across five countries and involved 43 000 nurses from the USA, Canada, England, Scotland and Germany. A consortium developed a nurse questionnaire dealing with issues related to nurses' perceptions of their working environments and quality of nursing care, job satisfaction, career plans, and feelings of job burnout. Their study found low morale, job dissatisfaction, burnout and intent to leave their current employers were common across the sample. Furthermore, nurses reported an increase in workload and non-nursing tasks, resulting in a decreased ability to complete nursing duties due to the competition created between nursing and non-nursing tasks. This context has the potential to contribute significantly to a decrease in the provision of quality care leading to negative outcomes for the patient and their family. A limitation to the study was the inclusion of only developed countries and sole reliance on self-report measures.

An additional online survey was posted by the ANA (2001b) on the health and safety of RNs in their work environment. A total of 4826 nurses from the USA responded. Quantitative data showed that the respondents' major health concern was the acute/chronic effects of stress and overwork (71%) they experienced. A contributing factor reported in the study was the perceived lack of safety in the work environment experienced by 80% of the RNs. Although the study was online and therefore likely reflects a motivated group of nurses, it did represent a cross-section

of nurses from every age group, years of experience, and types of care facilities.

Wickett et al.'s (2003) commentary on Janiszewski Goodin's (2003) article relating to the Australian perspective on the nursing shortage in Australia reported that Australian nurses also experienced additional pressures resulting from increased patient acuity combined with shorter hospital stays, similarly reported by Janiszewski Goodin (2003). Wickett et al.'s (2003) commentary noted that the substitution of RNs for unskilled workers as a cost-cutting measure has resulted in an increase in the workload and dissatisfaction of RNs, and they are playing a greater supervisory role rather than being direct caregivers. They also argue that removal of the RN from direct care might be a contributing factor to increased job dissatisfaction and therefore exacerbated stress levels (Wickett et al., 2003). Third, professional issues such as the perceived disparity between expectations of standards of care and what they are unable to sustain, and lack of career opportunities have also impacted on the recruitment and retention of nurses.

Buchanan and Considine (2002) conducted an Australian qualitative focus group study and found that the increased turnover and acuity of patients without adjusting staffing levels was a great source of stress for RNs. Other stressors identified by the study included: greater responsibility and roles; performing duties other than their speciality due to lack of staff; new technologies and teaching others to use them; and increased workload due to greater requirements of record-keeping and data collection. These factors have resulted in the nature of nursing being more stressful, less satisfying and less rewarding. Many nurses felt that these demands made them less able to cope with their role, leading to increased staff turnover and intention to leave. These findings support other Australian survey research previously conducted (Healy & McKay, 2000) that also found work overload to be a major source of stress for nurses. The sample were 129 RNs who volunteered to answer five standardized questionnaires as well as open-ended questions. Generalizability of the findings is limited by the moderate sample size and self-report nature of the study.

AGE AND ROLE STRESS

In addition to the problems of undersupply of nursing staff, the nursing population is experiencing an increasing average age, with an average age of 44 years in the USA, and forecast to have more than 40% older than 50 years in 10 years time (Janiszewski Goodin, 2003). The average age is similar in Australia and the UK (Mullen, 2003; Wickett et al., 2003). The implications of this are that most experienced nurses, mainly middle-aged women, will be leaving the profession at a time when supply is at its lowest (Janiszewski Goodin, 2003).

Santos et al. (2003) conducted a research study by administering the Occupational Stress Inventory - Revised Edition (OSI-R) to 694 nurses from four hospitals in the USA to determine the influence of age on the following aspects of role stress: job role overload, role insufficiency, role ambiguity, role boundary, responsibility, and physical environment. The OSI-R also assesses four types of personal strain: vocational, psychological, interpersonal and physical. The researchers conducted focus groups to clarify findings of the survey. The greatest problems identified were the physical demands and responsibility of inpatient nursing. Role boundary and role overload were key types of stress-inducing roles. Because of increased patient acuity and early discharge, the physical impact of treating sicker patients who require more intense care and treatments is magnified. All of the issues identified seemed to be amplified for the 'baby boomer' generation of RNs (those born between 1946 and 1964). Implications of these findings are discussed below in Strategies to reduce role stress. The findings of this study are limited by the fact that data were collected from

only one region and the voluntary nature of participation, leading to possible self-selection bias.

Results of a recent quantitative Japanese study of workplace stressors in hospital nurses (Lambert et al., 2004) differed to the research findings above. Role stress was defined as per the current article. The authors found that older age and more years worked as a nurse were negatively related to the stressors of workload, death/dying, inadequate preparation and uncertainty about treatment. These findings may be explained in terms of differing practices in Japan compared with Western nursing. In Japan, the younger, less experienced members of the workforce tend to be assigned an extra workload and more unpleasant aspects of nursing (Lambert et al., 2004). This study found that older nurses tended to distance themselves from others in order to cope with work stressors. The researchers also found a significant positive correlation between the likelihood of leaving the current nursing position and workplace stressors, workload, conflict with physicians, conflict with other nurses, lack of support, inadequate preparation and uncertainty about treatment. However, it is questionable whether the findings can be extrapolated beyond Japanese culture. The study findings are further limited by the large number of correlations used in this study that reduce statistical power.

ROLE STRESS AND NEW GRADUATES

Given the increased average age of the RN and the issues related to retention of the existing workforce, the sustainability of our workforce is in jeopardy. For our 'young' to be already exposed to such role stress indicates little resolution of the underlying issues. In addition to the previously mentioned stresses experienced by RNs, the new graduate has other pressures to contend with simultaneously. These may include a lack of confidence, unrealistic expectations by clinical staff, role conflict and role ambiguity, value conflicts, and lack of support (Kelly, 1998). Chang and Hancock's (2003) research study on role stress (defined as per the current article) among new graduates found that they experienced significant role ambiguity that was related to their job dissatisfaction over their initial 12 months. Transition to their new role as a RN was associated with a lack of clear and consistent information about the role. Additionally, conflicting role expectations were demanded of them. However, the findings of this study are limited by the sole reliance on self-report measures and only apply to new graduates.

ROLE STRESS AND MODES OF ORGANIZING WORK

The recurrent or unresolved problem of the need to retain qualified and experienced nurses highlights the need to study various aspects of nurses' work and environment, in which those factors affecting job satisfaction and aspects which contribute to the delivery of quality of care are exposed and acted upon. Makinen et al. (2003) conducted a research study investigating the relationship between modes of organizing nursing work and stressful characteristics of work. In hospital settings the three main modes of organizing nursing work are functional nursing, team nursing and primary nursing. Functional nursing is task-oriented with ward routine rather than individual patient needs being the priority (Makinen et al., 2003). Team nursing involves carrying out sets of activities in teams, while primary nursing involves patients being allocated to individual nurses (Makinen et al., 2003). However, nurses do not always work according to these three modes (Makinen et al., 2003). Modular nursing is another category, which is a modification of both team and primary nursing. Stressful job characteristics were measured by the

Occupational Stressor Questionnaire (Kivimaki & Lindstrom, 1995). This includes aspects such as work overload, high responsibility, and problems in interpersonal relations. The study found that the organization of nursing care was not strongly associated with stressful job characteristics. Interpersonal relationship problems such as communication problems among staff were less likely when patient-focused nursing modes were used, with this mode possibly reducing role ambiguity (Makinen et al., 2003). Nursing modes that provided opportunities for writing nursing notes were related to lower stress. The limitations of this study are that the findings cannot be generalized beyond medical and surgical wards to other specialties.

VIOLENCE

An emerging issue increasingly regarded as a source of role stress in nurses is violence. Nurses may receive violent acts from patients, relatives, other nurses and other professional groups (Jackson et al., 2002). International studies reported on from the USA (Kaye, 1996; Dalphond et al., 2000; Erickson & Williams-Evans, 2000), Sweden (Arnetz & Arnetz, 2000), and Australia (Fisher et al., 1995; Farrell, 1999; O'Connell et al., 2000) indicate that violence and harassment in the workplace are significant problems being confronted by nurses daily. Subsequently, nurses are reportedly at greater risk of assault in the workplace than other health professionals (Carter, 2000). Jackson et al. (2002), in their review of workplace violence and nurses, argue that the potential to be exposed to such aggression results in anxiety at work and may be related to sick leave, burnout and poor recruitment and retention rates. In their discussion paper, Wickett et al. (2003) argue that an increase in behavioral problems in Australian society is reflected in rising admissions of such individuals to the acute care setting. For instance, nurses are often involved in dealing with victims and perpetrators of violence (Jackson et al., 2002).

It was once the norm that acute hospitalization would entail a week's hospital stay, with patients highly dependent on nurses for care. Modern nursing care is more limited and patients are encouraged to be independent soon after treatment interventions. However, Wickett et al. (2003) discuss how some patients have the expectation that nurses should be providing all of their care, which can result in physical and verbal violence due to the mismatch in expectation. This situation adds to the role stress experienced by nurses and is one factor purported to be related to nurses leaving the profession (Wickett et al., 2003). The Royal College of Nursing (RCN) found similar issues to those in 1998 for nurses leaving the profession, but with the additional reasons of bullying and violence at work (RCN, 2000).

STRATEGIES TO REDUCE ROLE STRESS

Previous studies on role stress in nursing suggest that a range of strategies could be implemented to assist nurses with this experience. A qualitative study using grounded theory methodology found that the stress experienced by perioperative nurse managers was reduced through the use of hospital resources and peer support, referring to postmanagement education and information obtained from attending conferences (Schroeder & Worrall-Carter, 2002). In addition, they used team-building strategies, balanced priorities and engaged in social activities. While the generalizability of these findings are limited by their qualitative nature, further evidence that social support may assist in coping with role stress comes from research by Garrett and McDaniel (2001) who used a cross-sectional survey and found that the social climate of the workplace was negatively associated with burnout.

They suggest that ‘social networks are important during times of change and uncertainty in the work environment: a supportive workplace can protect against burnout.’ (p. 2). Others have concluded from their research that burnout may be prevented by fostering social relations in the organization (Janssen et al., 1999).

Flexibility in work hours may reduce role stress. Traditionally, nursing was seen as a vocation and women did not combine this career with parenting. Today, however, nursing remains a predominantly female workforce, but with many having dependant children. One of the major personal issues identified in Australia as influencing nursing retention is the lack of a ‘family friendly’ workplace. The strain of participating in a 7 day per week/24 h coverage roster reduces the resources available for nurses to cope with the demands of family roles. A research study of female nurses, combining partner and parent roles while working a rotating-shift roster, found that the interaction between work and family roles resulted in chronic fatigue that was a risk factor, especially when combined with the acute fatigue associated with shiftwork (Clissold et al., 2002). The researchers concluded that the need for flexible rostering is paramount in the retention of nurses. In terms of dealing with violence, Jackson et al. (2002) suggest that ‘effective protocols are needed to deal with bullying and abuse of power by nurses and others that are aimed at supporting nursing personnel’ (p. 19). They recommend that both employers and educational institutions ensure nurses are prepared to deal with acts of violence as they arise, and that employers demonstrate that they are committed to ensuring the safety of nurses. They believe that nurses need to feel that they are supported by their employer when reporting incidents of violence against them. However, evidence for the efficacy of these strategies is needed.

Concerning new graduates, Chang and Hancock (2003) recommend from their research findings that nurse unit managers (NUMs) should not only assess the impact of levels of role conflict and ambiguity on nurses’ adjustment to their role, but the impact of these factors on new graduates’ working environment. Adjustment could also be enhanced by the NUMs specifying the roles of the neophyte to both the individual and other staff. Recruitment, retention and facilitation of transition for new graduates can be enhanced by orientation, preceptorship and/or mentoring programs (Allanach & Jennings, 1990; Anderson, 1993; Boyle et al., 1996; Brans, 1997; Oermann & Moffitt-Wolf, 1997; Prebble & McDonald, 1997). Greenwood (2000) reported that factors such as formal unit orientation programs; a unit ‘climate’ of open communication; preceptor programs; the timely provision of constructive feedback; assignment congruence (tasks commensurate with competence) participative, democratic governance; appropriate advice and guidance from senior staff; and continuing staff development opportunities greatly assists new graduates in their experience of role stress and ambiguity. However, evidence for the relevance and appropriateness needs to be undertaken to inform practice. A review of the literature on stress management interventions in mental health nurses conducted by Edwards and Burnard (2003), that included research from 1966 to 2000, suggested that the most effective way to manage stress is a multilevel approach. First, to minimize or eliminate the stressors themselves, the organizational environment must be proactive rather than reactive in its management strategies. Due to the lack of research on the effectiveness of interventions it is difficult, however, to determine which specific techniques are the most effective. A second step is to use stress education and management strategies to reduce the negative effects of stress. Lastly, those who are experiencing the effects of stress should be assisted. Differing use of methodologies, tools and evaluation of different interventions, however, makes it difficult to compare the studies and generalize findings.

Although Edwards and Burnard (2003) looked specifically at mental health and

stress rather than role stress specifically, these recommendations are transferable to the context of role stress/strain being experienced in a general, hospital setting. To address the problem of an increasingly aging workforce and the finding that baby boomers experienced greater role stress (Santos et al., 2003), one possible strategy is to remove the mandatory age of retirement and encourage nurses to continue pursuing further study to prevent a drain of experience from the workforce. New and innovative work practices are needed to support the special needs of nurses in the older age bracket. These ideas, however, need to be tested for their utility. Workload is a dominant factor arising from this review as being related to role stress and strain. Given that Healy and McKay (2000) found that work stress, as measured by the Nursing Stress Scale (Gray-Toft & Anderson, 1981), was the strongest predictor of mood disturbance in nurses in their research study, along with the findings of other researchers (ANA, 2001a, 2001b; Buchanan & Considine 2002; Janiszewski Goodin, 2003), interventions aimed at reducing the impact of environmental stressors such as workload and staffing are recommended to decrease their stress levels and encourage retention of staff. Janssen et al. (1999) also suggest that emotional exhaustion can be reduced by paying attention to workload. For instance, apart from employing more staff, workload can be spread more equally among units, or roster duty techniques can be applied so that there is a more adequate spread of activities during a certain timespan. Wickett et al. (2003) suggest that nurses need to be better prepared to work in an environment of increasingly sicker patients with greater turnover. Again, however, these suggestions need to be tested for their impact.

Research has found a relationship between work motivation and work content (Janssen et al., 1999). A suggested strategy arising from this finding is to make elements of the job more challenging and worthwhile through varying tasks, providing more opportunities for autonomy and timely feedback. Job posting, job rotation, career counselling, career development and more comprehensive integrated human resource management programs can also assist with increasing work motivation, and thereby reducing the negative impact of role stress.

A recent theoretical article published by Lambert et al. (2003) presented strategies that could be used in the health care environment to enhance the three components of psychological hardiness (commitment, control and challenge), and thereby reduce the experience of stress in the workplace. Lambert et al. (2003) suggest that building commitment (defined as a sense of purpose and meaning expressed through active involvement in life's events [Lambert et al., 2003]) may be achieved through the following strategies: revising and rehearsing what you would do the next time the specific stressful event occurs; express yourself directly to the involved person(s); and rework the situation in your mind. Building control (tendency for the person to believe and act in a way that influences life's events rather than feeling helpless when confronted with adversity (Lambert et al., 2003) can be enhanced by seeking more information about the situation; trying to reduce stress (e.g. physical activity); trying to lighten or brighten the environment; or searching for a philosophical and/or spiritual meaning in the stressful experience. Building challenge (belief that change, instead of stability, is normal in life rather than a threat to security (Lambert et al., 2003), can be enhanced by use of interpersonal skills; searching for ways to keep a sense of perspective about the situation; broaden the range of influence and concern beyond the specific work situation; and cultivate an objective, intellectual attitude. Although the framework of psychological hardiness can be applied to role stress in nursing, it needs further empirical testing in terms of stress reduction outcomes.

In their research study of Japanese nurses, Lambert et al. (2004) found that there was a positive correlation between physical and mental health. Although causal

relationships cannot be demonstrated, their findings lend support for the hypothesis that an increase in workload affects the physical well-being of nurses. They also found that lack of support and the use of escape-avoidance as coping mechanisms were not helpful in terms of maintenance of mental health. This research suggests that contending with stressful events rather than avoiding them and the provision of support within the workplace are important ways of enhancing mental health and therefore coping with role stress.

The obvious strategy of reducing workload by employing more nursing staff is a stop-gap measure lacking sustainability due to the inadequate numbers of staff available to recruit from initially. As mentioned previously in this review, there exists a chronic short-age of nursing staff available to be recruited. Therefore this strategy is no longer considered a long-term resolution to the identified issue. Janiszewski Goodin (2003) proposes several strategies based on her review of the literature to increase retention and attract nurses to the workforce. These include improving personnel policies and benefits, career advancement opportunities, lifelong learning, flexible scheduling of work, and pay levels not fixed but commensurate with levels of preparation, experience, responsibility and performance. Janiszewski Goodin (2003) argues that 'only when the labor markets begin to respond to the imbalances of supply and demand in the nursing workforce will overall improvements in RN wages be seen' (p. 340).

Although the state of nursing role stress and the undersupply of the workforce in Australia reflects those of the USA, there are some signs of a turnaround. Nursing shortages are a significant problem in Australia, yet last year there was a sharp increase in the number of enrollments into Australian undergraduate nursing programs (Wickett et al., 2003). Wickett et al. (2003) attribute this to 'the concerted and collective efforts that have been made by the nursing profession in our country, and through government-funded media programmes, to recruit students into nursing programmes and to raise the profile of nursing in the wider community' (p. 344). Whether or not retention of these numbers can be maintained postgraduation, is yet to be determined.

CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH

This review has identified multiple factors related to the experience of role stress in nurses. Role stress, in particular, work overload, has been reported as one of the main reasons for nurses leaving the workforce. It is therefore a priority to not only reduce workload by attracting more nursing staff into the workforce, but to find new and innovative ways of supporting nurses in their experience of role stress. Some examples discussed in this article include use of stress education and management strategies to reduce the negative effects of stress; use of team-building strategies, balancing priorities, enhancing social support through engaging in social activities and peer support; flexibility in work hours; protocols to deal with violence; strategies to build commitment, control and challenge in the workplace; making elements of the job more challenging and worthwhile through varying tasks, providing more opportunities for autonomy and timely feedback; removing the mandatory age of retirement and encouraging nurses to continue pursuing further study to prevent a drain of experience from the workforce; increase nursing enrollments and use of strategies to attract nurses to the workforce. The next step is to empirically evaluate the efficacy of these strategies in reducing role stress in nurses.

One of the problems in comparing research on role stress/strain in nurses is the heterogeneity in the measures and methods used, making it difficult to compare findings. Furthermore, most studies rely on self-report measures only. It is

recommended that cross-cultural research comparing the experience of role stress in various countries using the same methods and measures be conducted. The authors are part of an international research team currently participating in such research. The research will enable an investigation of demographic, workplace stressors and coping mechanisms used as determinants of physical and mental health, similar to that performed by Lambert et al. (2004). This will assist in determining whether different countries are grappling with similar problems with regard to role stress, and whether differences lie in the levels of intensity or importance of various contributing factors. The study will also assist in predicting which ways of coping with role stress are most adaptive for RNs from an international perspective and specifically for each country studied.

REFERENCES

- Aiken L, Clarke S, Sloane D et al. Nurses' reports on hospital care in five countries. *Health Aff.* 2001; **20**: 43–53.
- Allenach B, Jennings B. Evaluating the effects of a nurse preceptorship programme. *JAN* 1990; **15**: 22–28.
- American Nurses Association (ANA). Analysis of American Nurses Association Staffing Survey. [Cited 9 September 2003]. Available from URL: <http://www.nursingworld.org/staffing> 2001a.
- American Nurses Association (ANA). Health and Safety Survey. [Cited 9 September 2003]. Available from URL: <http://www.nursingworld.org/about/mailroom.htm> 2001b.
- Andersson E. The perspective of student nurses and their perceptions of professional nursing during the nurse training programme. *J. Adv. Nurs.* 1993; **18**: 808–815.
- Arnetz J, Arnetz B. Implementation and evaluation of a practical intervention program for dealing with violence towards health care workers. *J. Adv. Nurs.* 2000; **31**: 668–680.
- Avallone I, Gibbon B. Nurses' perceptions of their work environment in a Nursing Development Unit. *J. Adv. Nurs.* 1998; **27**: 1193–1201.
- Boswell C. Work stress and job satisfaction for the community health nurse. *J. Community Health Nurs.* 1992; **9**: 221–227.
- Boyle D, Popkess-Vawter S, Taunton R. Socialization of new graduate nurses in critical care. *Heart Lung* 1996; **25**: 141–154.
- Brans L. Hit the Ground Running: Conference Report. Canberra: Royal College of Nursing, 1997.
- Buchanan J, Considine G. Stop Telling Us to Cope! A report for the NSW Nurses Association. Sydney: Australian Centre for Industrial Relations Research and Training (ACIRRT), University of Sydney, 2002.
- Buerhaus P. \$1 billion needed to address US shortage of nurses. *Aust. Nurs. J.* 2004; **11**: 21–26.
- Carter R. High risk of violence against nurses. *Nurs. Manag.* 2000; **6**: 5.
- Chang E, Hancock K. Role stress and role ambiguity in new nursing graduates in Australia. *Nurs. Health Sci.* 2003; **5**: 155–163.
- Chapman J. Collegial support linked to reduction of job stress. *Nurs. Manag.* 1993; **24**: 52–54.
- Cheng Y, Kawachi I, Coakley E, Schwartz ZJ, Colditz G. Association between psychosocial work characteristics and health functioning in American women. *BMJ* 2000; **320**: 1432–1436.
- Chung M, Corbett J. The burnout of nursing staff working with challenging behavior clients in hospital-based budgets and a community unit. *Int. J. Nurs. Stud.* 1998; **35**: 56–64.
- Clissold G, Smith P, Accutt B, Di Milia L. A study of female nurses combining partner and parent roles with working a continuous three-shift roster: The impact of sleep, fatigue and stress. *Contemp. Nurs.* 2002; **12**: 294–302.
- Cooper C. Introduction. In: Cooper C (ed). *Theories of Organizational Stress*. Oxford: Oxford University Press, 1998; 1–5.
- Dalphond D, Gessner M, Giblin E, Hijazzi K, Love C. Violence against emergency nurses. *J.*

- Emerg. Nurs. 2000; **26**: 105.
- Decker F. Occupational and non-occupational factors in jobsatisfaction and psychological distress among nurses. *Res. Nurs. Health* 1997; **20**: 453–464.
- Edwards D, Burnard P. A systematic review of stress and stress management interventions for mental health nurses. *J. Adv. Nurs.* 2003; **42**: 169–200.
- Erickson J, Williams-Evans S. Attitudes of emergency nurses regarding patient assaults. *J. Emerg. Nurs.* 2000; **26**: 210–215.
- Farrell G. Aggression in clinical settings: Nurses' views—a follow-up study. *J. Adv. Nurs.* 1999; **29**: 532–541.
- Fisher J, Bradshaw J, Currie B et al. Context of Silence: Violence and the Remote Area Nurse. Bundaberg: Central Queensland University, 1995.
- Fong C. A longitudinal study of the relationships between overload, social support, burnout among nursing educators. *J. Nurs. Educ.* 1993; **32**: 24–29.
- Foxall M, Zimmerman L, Standley R, Bene B. A comparison of frequency and sources of nursing job stress perceived by intensive care, hospice and medical-surgical nurses. *J. Adv. Nurs.* 1990; **15**: 577–584.
- Frisch S, Dembeck P, Shannon V. The head nurse: Perceptions of stress and ways of coping. *Can. J. Nurs. Adm.* 1991; **4**: 9–13.
- Garrett D, McDaniel A. A new look at nurse burnout: The effects of environmental uncertainty and social climate. *J. Nurs. Admin.* 2001; **31**: 91–96.
- Glass D, McKnight J, Valdimarsdottir J. Depression, burnout and perceptions of control in hospital nurses. *J. Consult. Clin. Psychol.* 1993; **61**: 147–155.
- Gray-Toft P, Anderson J. The Nursing Stress Scale: Development of an instrument. *J. Behav. Assess.* 1981; **3**: 11–23.
- Greenwood J. Critique of the graduate nurse: An international perspective. *Nurse Educ. Today* 2000; **20**: 17–23.
- Hatcher S, Laschinger H. Staff nurses' perceptions of job empowerment and level of burnout: A test of Kanter's theory of structural power in organizations. *Can. J. Nurs. Adm.* 1996; **9**: 74–94.
- Healy C, McKay M. Nursing stress: The effects of coping strategies and job satisfaction in a sample of Australian nurses. *J. Adv. Nurs.* 2000; **31**: 681–688.
- Jackson D, Clare J, Mannix J. Who would want to be a nurse? Violence in the workplace—a factor in recruitment and retention. *J. Nurs. Manage.* 2002; **10**: 13–20.
- Janiszewski Goodin H. The nursing shortage in the United States of America: An integrative review of the literature. *J. Adv. Nurs.* 2003; **43**: 335–350.
- Janssen P, de Jonge J, Bakker A. Specific determinants of intrinsic work motivation, burnout and turnover intentions: A study among nurses. *J. Adv. Nurs.* 1999; **29**: 1360–1369.
- Jex S, Beehr T. Emerging theoretical and methodological issues in the study of work related stress. *Res. Personnel Human Resources Manag.* 1991; **9**: 311–365.
- Kaye J. Sexual harassment and hostile environments in the perioperative area. *AORN J.* 1996; **63**: 443–449.
- Kelly B. Preserving moral integrity: A follow-up study of new graduate nurses. *J. Adv. Nurs.* 1998; **28**: 1134–1145.
- Kivimaki M, Lindstrom K. Effects of private self-consciousness and control on the occupational stress-strain relationship. *Stress Medicine* 1995; **11**: 7–16.
- Lally I, Pearce J. Intensive care nurses' perceptions of stress. *Nurs. Crit. Care* 1996; **1**: 17–25.
- Lambert V, Lambert C. Literature review of role stress/strain on nurses: An international perspective. *Nurs. Health Sci.* 2001; **3**: 161–172.
- Lambert V, Lambert C, Ito M. Workplace stressors, ways of coping and demographic characteristics as predictors of physical and emotional health of Japanese hospital nurses. *Int. J. Nurs. Stud.* 2004; **41**: 85–97.
- Lambert V, Lambert C, Yamase H. Psychological hardiness, workplace stress and related stress reduction strategies. *Nurs. Health Sci.* 2003; **5**: 181–184.
- Lazarus R, Folkman S. *Stress, Appraisal and Coping*. New York: Springer, 1984.
- Lee V, Henderson M. Occupational stress and organizational commitment in nurse administrators. *J. Nurs. Adm.* 1996; **26**: 21–28.

28 *Work Engagement And Occupational Stress In Nurses And Other Healthcare Workers: The Role Of Organizational And Personal Factors*

- Makinen A, Kivimakim M, Elovainio M, Virtanen M. Organization of nursing care and stressful work characteristics. *Nurs. Health Care Manag. Issues* 2003; **43**: 197–205.
- Manderino M, Brown M, Peters M, Wirtz R. Sources of stress for nurse practitioners. *J. Am. Acad. Nurs. Prac.* 1994; **6**: 155–159.
- Maslach C, Jackson S. MBI: Maslach Burnout Inventory, manual research edition. Palo Alto, CA: University of California, 1986.
- Mullen C. Commentary: An English perspective. *J. Adv. Nurs.* 2003; **43**: 345–348.
- Murray T. From outside the walls: A qualitative study of nurses who recently changed from hospital-based practice to home health care nursing. *J. Contin. Educ. Nurs.* 1998; **29**: 55–60.
- O’Connell B, Young J, Brooks J, Hutchings J, Lofthouse J. Nurses’ perceptions of the nature and frequency of aggression in general ward settings and high dependency areas. *J. Clin. Nurs.* 2000; **9**: 602–610.
- Oermann M, Moffitt-Wolf A. New graduates’ perceptions of clinical practice. *J. Cont. Ed. Nurs.* 1997; **28**: 20–25.
- Prebble K, McDonald B. Adaptation to the mental health setting: The lived experience of comprehensive graduates. *Aust. NZ. J. Ment. Health Nurs.* 1997; **6**: 30–36.
- RMIT University. Newsletter. Melbourne: Department of Nursing and Midwifery, Business Unit, Life Sciences. RMIT University, 2003.
- Robinson J, Lewis D. Coping with ICU work-related stressors: A study. *Crit. Care Nurs.* 1990; **14**: 32–35.
- Royal College of Nursing (RCN). Local Needs, Local Solutions. London: Royal College of Nursing (UK), 2000.
- Santos S, Carrol C, Cox K et al. Baby boomer nurses bearing the burden of care: A four site study of stress, strain and coping for impatient registered nurses. *J. Nurs. Adm.* 2003; **33**: 243–250.